

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 :

C12N 9/54

A1

(11) International Publication Number:

WO 96/34946

(43) International Publication Date:

7 November 1996 (07.11.96)

(21) International Application Number: PCT/DK96/00207

(22) International Filing Date: 2 May 1996 (02.05.96)

(30) Priority Data:

0519/95

5 May 1995 (05.05.95)

DK

0421/96

12 April 1996 (12.04.96)

DK

(71) Applicant: NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK).

(72) Inventors: SIERKSTRA, L., N.; Unilever N.V., Weena 455, NL-3013 AL Rotterdam (NL). KLUGKIST, J.; Unilever N.V., Weena 455, NL-3013 AL Rotterdam (NL). MARKVARSEN, Peter; Novo Nordisk a/s, Novo Allé, DK-2880 Bagsværd (DK). VON DER OSTEN, Claus; Novo Nordisk a/s, Novo Allé, DK-2880 Bagsværd (DK). BAUDITZ, Peter; Novo Nordisk a/s, Novo Allé, DK-2880 Bagsværd (DK).

(74) Common Representative: NOVO NORDISK A/S; Corporate Patents, Novo Allé, DK-2880 Bagsværd (DK).

(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

With international search report.

(54) Title: PROTEASE VARIANTS AND COMPOSITIONS

(57) Abstract

A subtilisin having improved storage stability or improved performance in detergents, wherein one or more amino acid residues situated in, or in the vicinity of a hydrophobic domain of the parent subtilase have been substituted for an amino acid residue more hydrophobic than the original residue, said hydrophobic domain comprising the residues corresponding to residues P129, P131, I165, Y167, Y171 or BLS309 (in BASBPN numbering), and said residues in the vicinity thereof comprises residues corresponding to the residues E136, G159, S164, R170, A194, and G195 of BLS309 (in BASBPN numbering), with the exception of the R170M, R170I and R170Y variant of BLS309.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Title: Protease Variants and Compositions

TECHNICAL FIELD

5 This invention relates to novel mutant enzymes or enzyme variants useful in formulating detergent compositions and exhibiting improved storage stability while retaining or improving their wash performance; cleaning and detergent compositions containing said enzymes; mutated genes coding for
10 the expression of said enzymes when inserted into a suitable host cell or organism; and such host cells transformed therewith and capable of expressing said enzyme variants.

15

BACKGROUND OF THE INVENTION

In the detergent industry enzymes have for more than 30 years been implemented in washing formulations. Enzymes used in such formulations comprise proteases, lipases, amylases, cellulases,
20 as well as other enzymes, or mixtures thereof. Commercially most important are proteases.

Although proteases have been used in the detergent industry for more than 30 years, much remains unknown as to details of how
25 these enzymes interact with substrates and/or other substances present in e.g. detergent compositions. Some factors related to specific residues and influencing certain properties, such as oxidative and thermal stability in general have been elucidated, but much remains to be found out. Also, it is still
30 not exactly known which physical or chemical characteristics are responsible for a good washing performance or stability of a protease in a specific detergent composition.

SUBSTITUTE SHEET

The currently used proteases have for the most part been found by isolating proteases from nature and testing them in detergent formulations.

5

An increasing number of commercially used protease are protein engineered variants of the corresponding naturally occurring wild type protease, e.g. DURAZYM® (Novo Nordisk A/S), RELEASE® (Novo Nordisk A/S), MAXAPEM® (Gist-Brocades N.V.), PURAFECT®
10 (Genencor International, Inc.).

Therefore, an object of the present invention, is to provide improved protein engineered protease variants, especially for use in the detergent industry.

15

PROTEASES

Enzymes cleaving the amide linkages in protein substrates are classified as proteases, or (interchangeably) peptidases (see Walsh, 1979, *Enzymatic Reaction Mechanisms*. W.H. Freeman and
20 Company, San Francisco, Chapter 3). Bacteria of the *Bacillus* species secrete two extracellular species of protease, a neutral, or metalloprotease, and an alkaline protease which is functionally a serine endopeptidase and usually referred to as subtilisin. Secretion of these proteases has been linked to the
25 bacterial growth cycle, with greatest expression of protease during the stationary phase, when sporulation also occurs. Joliffe et al. (1980) *J. Bacteriol* 141 1199-1208, have suggested that *Bacillus* proteases function in cell wall turnover.

30

SUBTILASES

A serine protease is an enzyme which catalyzes the hydrolysis of peptide bonds, and in which there is an essential serine residue at the active site (White, Handler and Smith, 1973

SUBSTITUTE SHEET

"Principles of Biochemistry," Fifth Edition, McGraw-Hill Book Company, NY, pp. 271-272).

The bacterial serine proteases have molecular weights in the 5 20,000 to 45,000 Daltons range. They are inhibited by diisopropylfluorophosphate. They hydrolyze simple terminal esters and are similar in activity to eukaryotic chymotrypsin, also a serine protease. A more narrow term, alkaline protease, covering a sub-group, reflects the high pH optimum of some of 10 the serine proteases, from pH 9.0 to 11.0 (for review, see Priest (1977) *Bacteriological Rev.* 41 711-753).

A sub-group of the serine proteases tentatively designated subtilases has been proposed by Siezen et al., *Protein Engng.* 4 15 (1991) 719-737. They are defined by homology analysis of more than 40 amino acid sequences of serine proteases previously referred to as subtilisin-like proteases. A subtilisin was previously defined as a serine protease produced by Gram-positive bacteria or fungi, and according to Siezen et al. now 20 is a subgroup of the subtilases. A wide variety of subtilisins have been identified, and the amino acid sequence of a number of subtilisins have been determined. These include more than six subtilisins from *Bacillus* strains, namely, subtilisin 168, subtilisin BPN', subtilisin Carlsberg, subtilisin Y, 25 subtilisin amylosacchariticus, and mesentericopeptidase (Kurihara et al. (1972) *J. Biol. Chem.* 247 5629-5631; Wells et al. (1983) *Nucleic Acids Res.* 11 7911-7925; Stahl and Ferrari (1984) *J. Bacteriol.* 159 811-819, Jacobs et al. (1985) *Nucl. Acids Res.* 13 8913-8926; Nedkov et al. (1985) *Biol. Chem.* 30 Hoppe-Seyler 366 421-430, Svendsen et al. (1986) *FEBS Lett.* 196 228-232), one subtilisin from an actinomycetales, thermitase from *Thermoactinomyces vulgaris* (Meloun et al. (1985) *FEBS Lett.* 198 195-200), and one fungal subtilisin, proteinase K from *Tritirachium album* (Jany and Mayer (1985) *Biol. Chem.*

SUBSTITUTE SHEET

Hoppe-Seyler 366 584-492). for further reference Table I from Siezen et al. has been reproduced below.

Subtilisins are well-characterized physically and chemically. In addition to knowledge of the primary structure (amino acid sequence) of these enzymes, over 50 high resolution X-ray structures of subtilisins have been determined which delineate the binding of substrate, transition state, products, at least three different protease inhibitors, and define the structural consequences for natural variation (Kraut (1977) Ann. Rev. Biochem. 46 331-358).

In the context of this application substrate should be interpreted in its broadest form as comprising a compound containing at least one peptide bond susceptible to hydrolysis by a subtilisin protease.

Also the expression "product" should in the context of this invention be interpreted to include the products of a hydrolysis reaction involving a subtilisin protease. A product may be the substrate in a subsequent hydrolysis reaction.

One subgroup of the subtilases, I-S1, comprises the "classical" subtilisins, such as subtilisin 168, subtilisin BPN', subtilisin Carlsberg (ALCALASE®, NOVO NORDISK A/S), and subtilisin DY.

A further subgroup of the subtilases I-S2, is recognised by Siezen et al. (*supra*). Sub-group I-S2 proteases are described as highly alkaline subtilisins and comprise enzymes such as subtilisin PB92 (MAXACAL®, Gist-Brocades NV), subtilisin 309 (SAVINASE®, NOVO NORDISK A/S), subtilisin 147 (ESPERASE®, NOVO NORDISK A/S), and alkaline elastase YaB.

SUBSTITUTE SHEET

In the context of this invention, a subtilase variant or mutated subtilase means a subtilase that has been produced by an organism which is expressing a mutant gene derived from a parent microorganism which possessed an original or parent gene and which produced a corresponding parent enzyme, the parent gene having been mutated in order to produce the mutant gene from which said mutated subtilisin protease is produced when expressed in a suitable host.

- 10 Random and site-directed mutations of the subtilase gene have both arisen from knowledge of the physical and chemical properties of the enzyme and contributed information relating to subtilase's catalytic activity, substrate specificity, tertiary structure, etc. (Wells et al. (1987) *Proc. Natl. Acad. Sci. U.S.A.* 84; 1219-1223; Wells et al. (1986) *Phil. Trans. R. Soc. Lond.A.* 317 415-423; Hwang and Warshel (1987) *Biochem.* 26 2669-2673; Rao et al., (1987) *Nature* 328 551-554.

More recent publications covering this area are Carter et al. (1989) *Proteins* 6 240-248 relating to design of variants that cleave a specific target sequence in a substrate (positions 24 and 64); Graycar et al. (1992) *Annals of the New York Academy of Sciences* 672 71-79 discussing a number of previously published results; and Takagi (1993) *Int. J. Biochem.* 25 307-312 also reviewing previous results.

Especially site-directed mutagenesis of the subtilisin genes has attracted much attention, and various mutations are described in the following patent applications and patents:

30

EP publ. no. 130756 (GENENTECH) (corresponding to US Reissue Patent No. 34,606 (GENENCOR)) relating to site specific or randomly generated mutations in "carbonyl hydrolases" and subsequent screening of the mutated enzymes for various properties, such as k_{cat}/K_m ratio, pH-activity profile, and

SUBSTITUTE SHEET

oxidation stability. This publication reveals that site-specific mutation is feasible, and that mutation of subtilisin BPN' in certain specified positions, i.e. ⁻¹Tyr, ³²Asp, ¹⁵⁵Asn, ¹⁰⁴Tyr, ²²²Met, ¹⁶⁶Gly, ⁶⁴His, ¹⁶⁹Gly, ¹⁸⁹Phe, ³³Ser, ²²¹Ser, ²¹⁷Tyr, ¹⁵⁶Glu or ¹⁵²Ala, provide for enzymes exhibiting altered properties. Since these positions all except position -1 were known to be involved in the functioning of the enzyme prior to the filing of the application, and therefore evident to select, this application does not contribute much to solving the problem of deciding where to introduce mutations in order to obtain enzymes with desired properties.

EP publ. no. 214435 (HENKEL) relating to cloning and expression of subtilisin Carlsberg and two mutants thereof. In this application no reason for mutation of ¹⁵⁸Asp to ¹⁵⁸Ser and ¹⁶¹Ser to ¹⁶¹Asp is provided.

In International patent publication No. WO 87/04461 (AMGEN) it is proposed to reduce the number of Asn-Gly sequences present in the parent enzyme in order to obtain mutated enzymes exhibiting improved pH and heat stabilities, in the application emphasis is put on removing, mutating, or modifying the ¹⁰⁹Asn and the ²¹⁸Asn residues in subtilisin BPN'. No examples are provided for any deletions or for modifying the Gly-residues.

25

International patent publication No. WO 87/05050 (GENEX) discloses random mutation and subsequent screening of a large number of mutants of subtilisin BPN' for improved properties. In the application mutations are described in positions ²¹⁸Asn, ¹³¹Gly, ²⁵⁴Thr, ¹⁶⁶Gly, ¹¹⁶Ala, ¹⁸⁸Ser, ¹²⁶Leu, and ⁵³Ser.

In EP Publication no. 251 446 (GENENCOR) it is described how homology considerations at both primary and tertiary structural levels may be applied to identify equivalent amino acid residues whether conserved or not. This information together

SUBSTITUTE SHEET

with the inventors knowledge of the tertiary structure of subtilisin BPN' lead the inventors to select a number of positions susceptible to mutation with an expectation of obtaining mutants with altered properties. The positions so identified are: ¹²⁴Met, ²²²Met, ¹⁰⁴Tyr, ¹⁵²Ala, ¹⁵⁶Glu, ¹⁶⁶Gly, ¹⁶⁹Gly, ¹⁸⁹Phe, ²¹⁷Tyr. Also ¹⁵⁵Asn, ²¹Tyr, ²²Thr, ²⁴Ser, ³²Asp, ³³Ser, ³⁶Asp, ⁴⁶Gly, ⁴⁸Ala, ⁴⁹Ser, ⁵⁰Met, ⁷⁷Asn, ⁸⁷Ser, ⁹⁴Lys, ⁹⁵Val, ⁹⁶Leu, ¹⁰⁷Ile, ¹¹⁰Gly, ¹⁷⁰Lys, ¹⁷¹Tyr, ¹⁷²Pro, ¹⁹⁷Asp, ¹⁹⁹Met, ²⁰⁴Ser, ²¹³Lys, and ²²¹Ser, which positions are identified as being expected to influence various properties of the enzyme. Also, a number of mutations are exemplified to support these suggestions. In addition to single mutations in these positions the inventors also performed a number of multiple mutations. Further the inventors identify ²¹⁵Gly, ⁶⁷His, ¹²⁶Leu, ¹³⁵Leu, and amino acid residues within the segments 97-103, 126-129, 213-215, and 152-172 as having interest, but mutations in any of these positions are not exemplified.

Especially of interest for the purpose of the present invention the inventors of EP 251 446 suggest to substitute ¹⁷⁰Lys (in subtilisin BPN', type I-S1), specifically they suggest to introduce Glu or Arg for the original Lys. It appears that the Glu variant was produced and it was found that it was highly susceptible to autolytic degradation (cf. pages 48, 121, 123 (Table XXI includes an obvious error, but indicates a reduction in autolysis half-time from 86 to 13 minutes) and Fig. 32).

EP publ. no. 260105 (GENENCOR) describes modification of certain properties in enzymes containing a catalytic triad by selecting an amino acid residue within about 15 Å from the catalytic triad and replace the selected amino acid residue with another residue. Enzymes of the subtilase type described in the present specification are specifically mentioned as belonging to the class of enzymes containing a catalytic triad.

SUBSTITUTE SHEET

In subtilisins positions 222 and 217 are indicated as preferred positions for replacement.

Also, it has been shown by Thomas, Russell, and Fersht (1985) 5 *Nature* 318 375-376 that exchange of ⁹⁹Asp into ⁹⁹Ser in subtilisin BPN' changes the pH dependency of the enzyme.

In a subsequent article (1987) *J. Mol. Biol.* 193 803-813, the same authors also discuss the substitution of ¹⁵⁶Ser in place of 10 ¹⁵⁶Glu.

Both these mutations are within a distance of about 15Å from the active ⁶⁴His.

15 In *Nature* 328 496-500 (1987) Russel and Fersht discuss the results of their experiments and present rules for changing pH-activity profiles by mutating an enzyme to obtain changes in surface charge.

20 WO 88/08028 (Genex) and WO 88/08033 (Amgen) both relate to modifications of amino acid residues in the calcium binding sites of subtilisin BPN'. The enzyme is said to be stabilized by substituting more negatively charged residues for the original ones.

25

In WO 89/06279 (NOVO NORDISK A/S) position 170 is indicated as interesting and it is suggested to replace the existing residue with Tyr. However, no data are given in respect of such a variant. In WO 91/00345 (NOVO NORDISK A/S) the same suggestion 30 is made, and it is shown that the Tyr variant of position 170 in subtilisin 309 (type I-S2) exhibits an improved wash performance in detergents at a pH of about 8 (variant S003 in Tables III, IV, V, VI, VIII, X). The same substitution in combination with other substitutions in other positions also 35 indicates an improved wash performance(S004, S011-S014, S022-

SUBSTITUTE SHEET

S024, S019, S020, S203, S225, S227 in the same Table and Table VII) all in accordance with the generic concept of said application.

5 In EP 525 610 A1 (SOLVAY) it is suggested to improve the stability of the enzyme (a type I-S2 subtilase closely related to subtilisin PB92) towards ionic tensides by decreasing the hydrophobicity in certain surface regions thereof. It is consequently suggested to substitute Gln for the Arg in 10 position 164 (170 if using BPN' numbering). No variants comprising this substitution are disclosed in the application.

In WO 94/02618 (GIST-BROCADES N.V.) a number of position 164 (170 if using BPN' numbering) variants of the I-S2 type 15 subtilisin PB92 are described. Examples are provided showing substitution of Met, Val, Tyr, Ile, for the original Arg. Wash performance testing in powder detergents of the variants indicates a slight improvement. Especially for the Ile variant wash performance tests on cacao an improvement of about 20-30% 20 is indicated. No stability data are provided.

In WO 95/30011, WO 95/30010, and WO 95/29979 (PROCTER & GAMBLE COMPANY) describe 6 regions, especially position 199-220 (BPN' numbering), in both Subtilisin BPN' and subtilisin 309, which 25 are designed to change (i.e. decrease) the adsorption of the enzyme to surface-bound soils. It is suggested that decreased adsorption by an enzyme to a substrate results in better detergent cleaning performance. No specific detergent wash performance data are provided for the suggested variants.

30

WO 95/27049 (SOLVAY S.A.) describes a subtilisin 309 type protease with following mutations: N43R+N116R+N117R (BPN' numbering). Data indicate the corresponding variant is having improved stability, compared to wildtype.

35

SUBSTITUTE SHEET

INDUSTRIAL APPLICATIONS OF SUBTILASES

Proteases such as subtilisins have found much utility in industry, particularly in detergent formulations, as they are useful for removing proteinaceous stains.

At present at least the following proteases are known to be commercially available and many of them are marketed in large quantities in many countries of the world.

Subtilisin BPN' or Novo, available from e.g. SIGMA, St. Louis, U.S.A.

Subtilisin Carlsberg, marketed by NOVO NORDISK A/S (Denmark) as ALCALASE® and by Gist-Brocades N.V. (Holland) as MAXATASE®;

Both of these belong to subtilase subgroup I-S1

Among the subtilase sub-group I-S2 the following are known to be marketed.

A *Bacillus lentus* subtilisin, subtilisin 309, marketed by NOVO NORDISK A/S (Denmark) as SAVINASE®. A protein engineered variant of this enzyme is marketed as DURAZYM®.

Enzymes closely resembling SAVINASE®, such as subtilisin PB92, MAXACAL® marketed by Gist-Brocades N.V. (a protein engineered variant of this enzyme is marketed as MAXAPEM®), OPTICLEAN® marketed by SOLVAY et Cie. and PURAFECT® marketed by GENENCOR International.

A *Bacillus lentus* subtilisin, subtilisin 147, marketed by NOVO NORDISK A/S (Denmark) as ESPERASE®;

35

SUBSTITUTE SHEET

To be effective, however, such enzymes must not only exhibit activity under washing conditions, but must also be compatible with other detergent components during detergent production and storage.

5

For example, subtilisins may be used in combination with other enzymes active against other substrates, and the selected subtilisin should possess stability towards such enzymes, and also the selected subtilisin preferably should not catalyze
10 degradation of the other enzymes. Also, the chosen subtilisin should be resistant to the action from other components in the detergent formulation, such as bleaching agents, oxidizing agents, etc., in particular an enzyme to be used in a detergent formulation should be stable with respect to the oxidizing
15 power, calcium binding properties, and pH conditions rendered by the non-enzymatic components in the detergent during storage and in the wash liquor during wash.

The ability of an enzyme to catalyze the degradation of various
20 naturally occurring substrates present on the objects to be cleaned during e.g. wash is often referred to as its washing ability, washability, detergency, or wash performance. Throughout this application the term wash performance will be used to encompass this property.

25

The ability of an enzyme to remain active in the presence of other components of a detergent composition prior to being put to use (normally by adding water in the washing process) is usually referred to as storage stability or shelf life. It is
30 often measured as half-life, $t_{1/2}$. We will use the expression storage stability for this property throughout this application to encompass this property.

Naturally occurring subtilisins have been found to possess
35 properties which are highly variable in relation to their

SUBSTITUTE SHEET

washing power or ability under variations in parameters such as pH. Several of the above marketed detergent proteases, indeed, have a better performance than those marketed about 20 years ago, but for optimal performance each enzyme has its own 5 specific conditions regarding formulation and wash conditions, e.g. pH, temperature, ionic strength (=I), active system (tensides, surfactants, bleaching agent, etc.), builders, etc.

As a consequence it is found that an enzyme possessing 10 desirable properties at low pH and low I may be less attractive at more alkaline conditions and high I, or an enzyme exhibiting fine properties at high pH and high I may be less attractive at low pH, low I conditions.

15 Also, it has been found that the storage stability differs between the enzymes, but it has further been found that a specific enzyme exhibits large variations in storage stability in respect of different detergent formulations, dependent upon a number of parameters, such as pH, pI, bleach system, 20 tensides, etc., and upon the physical state of the detergent compositions, which may be in powder, dust, or liquid form. Furthermore it may be concentrated or dilute.

The advent and development of recombinant DNA techniques has 25 had a profound influence in the field of protein chemistry.

Through the application of this technology it is possible now to construct enzymes having desired amino acid sequences, and as indicated above a fair amount of research has been devoted 30 to designing subtilisins with altered properties.

Among the proposals the technique of producing and screening a large number of mutated enzymes as described in EP publ. no. 130756 (GENENTECH) (US Reissue Patent No. 34,606 (GENENCOR)) 35 and International patent publ. no. WO 87/05050 (GENEX)

SUBSTITUTE SHEET

correspond to a large extend to the classical method of isolating native enzymes, submit them to classical mutagenesis programs (using radiation or chemical mutagens) and screen them for their properties. The difference lies in that these methods are more efficient through the knowledge of the presence of a large number of variant enzymes substituted in a specific position.

A subtilisin enzyme typically comprises about 275 amino acid residues. Each residue is capable of being 1 out of 20 possible naturally occurring amino acids.

Therefore one very serious draw-back in that procedure is the very large number of mutations generated that have to be submitted to a number of preliminary screenings to determine their properties.

A procedure as outlined in these patent applications will consequently only be slightly better than the traditional random mutation procedures which have been known for years.

The other known techniques relate to changing specific properties, such as oxidation stability, thermal stability, Ca-stability, transesterification and hydrolysis rate (EP publ. no. 260105 (GENENCOR)), pH-activity profile (Thomas, Russell, and Fersht, *supra*), and substrate specificity (International patent publ. no. WO 88/07578 (GENENTECH)). None of these publications relates to changing either the wash performance of enzymes or their storage stability.

30

In International Patent Application no. PCT/DK 88/00002 (NOVO NORDISK A/S) it is proposed to use the concept of homology comparison to determine which amino acid positions should be selected for mutation and which amino acids should be

SUBSTITUTE SHEET

substituted in these positions in order to obtain a desired change in wash performance.

By using such a procedure the task of screening is reduced 5 drastically, since the number of mutants generated is much smaller, but with that procedure it is only foreseen that enzymes exhibiting the combined useful properties of the parent enzyme and the enzyme used in the comparison may be obtained.

10 Thus, as indicated above no relationship has yet been identified between well defined properties of an enzyme such as those mentioned above and the wash performance and storage stability of an enzyme in various detergent compositions.

15 The problem seems to be that although much research has been directed at revealing the mechanism of enzyme activity, still only little is known about the factors in structure and amino acid residue combination that determine the properties, such as storage stability in detergents, of enzymes in relation to most 20 of their characteristics, especially when the enzymes are present in complex mixtures.

Consequently there still exists a need for further improvement and tailoring of enzymes to detergent systems, as well as a 25 better understanding of the mechanism of protease action and degradation in the practical use of cleaning or detergent compositions. Such an understanding could result in rules which may be applied for selecting mutations that with a reasonable degree of certainty will result in an enzyme exhibiting 30 improved storage stability under specified conditions in a detergent composition.

SUBSTITUTE SHEET

SUMMARY OF THE INVENTION

5 It has now surprisingly been found that a subtilase variant having improved storage stability and/or improved performance in detergents, can be obtained by substituting one or more amino acid residues situated in, or in the vicinity of a hydrophobic domain of the parent subtilase for an amino acid
10 residue more hydrophobic than the original residue, said hydrophobic domain comprising the residues corresponding to residues P129, P131, I165, Y167, Y171 of BLS309 (in BASBPN numbering), and said residues in the vicinity thereof comprises residues corresponding to the residues E136, G159, S164, R170,
15 A194, and G195 of BLS309 (in BASBPN numbering), with the exception of the R170M, R170I and R170V variants of BABP92.

The present invention relates consequently in its first aspect to enzyme variants exhibiting improved stability and/or
20 improved wash performance in detergent.

In its second aspect the invention relates to DNA constructs capable of expressing the enzymes of the first aspect, when inserted in a suitable manner into a host cell that
25 subsequently is brought to express the subtilisin enzyme(s) of the first aspect.

In a third aspect the invention relates to the production of the subtilisin enzymes of the invention by inserting a DNA
30 construct according to the second aspect into a suitable host, cultivating the host to express the desired subtilase enzyme, and recovering the enzyme product.

SUBSTITUTE SHEET

The invention relates, in part, but is not limited to, mutants of the genes expressing the subtilase sub-group I-S2 enzymes and the ensuing enzyme variants, as indicated above.

Other subtilase gene variants encompassed by the invention are 5 such as those of the subtilase subgroup I-S1, e.g. Subtilisin BPN', and Subtilisin Carlsberg genes and ensuing variant Subtilisin BPN', Proteinase K, and Subtilisin Carlsberg enzymes, which exhibit improved stability in concentrated liquid detergents.

10

Still further subtilase gene variants encompassed by the invention are such as Proteinase K and other genes and ensuing variant Proteinase K, and other subtilase enzymes, which exhibit improved stability in concentrated liquid detergents.

15

Other examples of parent subtilase enzymes that can be modified in accordance with the invention are listed in Table I.

Further the invention relates to the use of the mutant enzymes 20 in cleaning compositions and cleaning compositions comprising the mutant enzymes, especially detergent compositions comprising the mutant subtilisin enzymes. Specifically the invention relates to concentrated liquid detergent compositions comprising such enzyme variants.

25

SUBSTITUTE SHEET

ABBREVIATIONSAMINO ACIDS

A	=	Ala	=	Alanine
5 V	=	Val	=	Valine
L	=	Leu	=	Leucine
I	=	Ile	=	Isoleucine
P	=	Pro	=	Proline
F	=	Phe	=	Phenylalanine
10 W	=	Trp	=	Tryptophan
M	=	Met	=	Methionine
G	=	Gly	=	Glycine
S	=	Ser	=	Serine
T	=	Thr	=	Threonine
15 C	=	Cys	=	Cysteine
Y	=	Tyr	=	Tyrosine
N	=	Asn	=	Asparagine
Q	=	Gln	=	Glutamine
D	=	Asp	=	Aspartic Acid
20 E	=	Glu	=	Glutamic Acid
K	=	Lys	=	Lysine
R	=	Arg	=	Arginine
H	=	His	=	Histidine
X	=	Xaa	=	Any amino acid

25 NUCLEIC ACID BASES

A	=	Adenine
G	=	Guanine
C	=	Cytosine
T	=	Thymine (only in DNA)
30 U	=	Uracil (only in RNA)

VARIANTS

In describing the various enzyme variants produced or contemplated according to the invention, the following nomenclatures have been adapted for ease of reference:

SUBSTITUTE SHEET

Original amino acid(s) position(s) substituted amino acid(s)

According to this the substitution of Glutamic acid for glycine 5 in position 195 is designated as:

Gly 195 Glu or G195E

a deletion of glycine in the same position is:

Gly 195 * or G195*

and insertion of an additional amino acid residue such as 10 lysine is:

Gly 195 GlyLys or G195GK

Where a deletion in comparison with the sequence used for the numbering is indicated, an insertion in such a position is 15 indicated as:

* 36 Asp or *36D

for insertion of an aspartic acid in position 36

Multiple mutations are separated by pluses, i.e.:

20 Arg 170 Tyr + Gly 195 Glu or R170Y+G195E

representing mutations in positions 170 and 195 substituting tyrosine and glutamic acid for arginine and glycine, respectively.

25

SUBSTITUTE SHEET

POSITIONS

In describing the variants in this application and in the appended claims use is made of the alignment of various subtilases in Siezen et al., *Supra*. In other publications relating to subtilases other alignments or the numbering of specific enzymes have been used. It is a routine matter for the skilled person to establish the position of a specific residue in the numbering used here. Reference is also made to Fig. 1 showing an alignment of residues relevant for the present invention from a large number of subtilases. Reference is also made to Table I of WO 91/00345 showing an alignment of residues relevant for the present invention from a number of subtilases.

15

TABLE I

Presently established Subtilases (from Siezen et al., *supra*)

Organism	cdNA, gene	enzyme	acronym
----------	---------------	--------	---------

PROKARYOTES20 Bacteria: Gram-positive

<i>Bacillus subtilis</i> 168	apr A	subtilisin I168, apr	
ABSS168			

<i>Bacillus amyloliquefaciens</i> apr		subtilisinBPN' (NOVO)	BASBPN
---------------------------------------	--	-----------------------	--------

<i>Bacillus subtilis</i> DY	-	subtilisin DY	BSSDY
-----------------------------	---	---------------	-------

25 <i>Bacillus licheniformis</i>	+	subtilisin Carlsberg	BLSCAR
----------------------------------	---	----------------------	--------

<i>Bacillus lentus</i>	+	subtilisin 147	BLS147
------------------------	---	----------------	--------

<i>Bacillus alcalophilus</i> PB92	+	subtilisin PB92	BAPB92
-----------------------------------	---	-----------------	--------

<i>Bacillus</i> sp. DSM 4828	-	alkaline protease	BDSM48
------------------------------	---	-------------------	--------

<i>Bacillus</i> YaB	ale	alkaline elastase YaB	BYSYAB
---------------------	-----	-----------------------	--------

30 <i>Bacillus subtilis</i> 168	epr	min. extracell. prot.	BSEPR
---------------------------------	-----	-----------------------	-------

<i>Bacillus subtilis</i>	bpf	bacillopeptidase F	BSBPF
--------------------------	-----	--------------------	-------

<i>Bacillus subtilis</i> IFO3013	ispl	intracell. ser. prot. 1	BSISP1
----------------------------------	------	-------------------------	--------

<i>Bacillus subtilis</i> A50	-	intracell. ser. prot.	BSIA50
------------------------------	---	-----------------------	--------

<i>Bacillus thuringiensis</i>	-	extracell. ser. prot.	BTFINI
-------------------------------	---	-----------------------	--------

SUBSTITUTE SHEET

	<i>Bacillus cereus</i>	-	extracell. ser. prot.	BCESPR
	<i>Nocardiopsis dassonvillei</i>	-	alkaline ser. prot.	NDAPII
	<i>Thermoactinomyces vulgaris</i>	-	thermitase	TVTHER
	<i>Enterococcus faecalis</i>	cylA	cytolysin component A	EFCYLA
5	<i>Staphylococcus epidermidis</i>	epiP	epidermin lead. prot.	SEEPiP
	<i>Streptococcus pyrogenes</i>	scpA	C5a peptidase	SPSCPA
	<i>Lactococcus lactis</i> SK11	prtP	SK11 cell wall prot.	LLSK11
	<u>Bacteria: Gram-negative</u>			
	<i>Dichelobacter nodosus</i>	+	basic protease	DNEBPR
10	<i>Xanthomonas campestris</i>	+	extracellular prot.	XCEXP
	<i>Serratia marcescens</i>	+	extracell. ser. prot.	SMEXSP
	<i>Thermus aquaticus</i> YT-1	pstI	aqualysin I	TAAQUA
	<i>Thermus</i> rT41A	+	T41A protease	TRT41A
	<i>Vibrio alginolyticus</i>	proA	protease A	VAPROA
15	<i>Streptomyces rutgersensis</i>	-	proteinase D	SRESPD
	<u>Archaea</u>			
	halophilic strain 172P1	-	halophil extra. prot.	ARB172
	<u>Cyanobacteria</u>			
	<i>Anabaena variabilis</i>	prcA	Ca-dependent protease	AVPRCA
20	<u>LOWER EUKARYOTES</u>			
	<u>Fungi</u>			
	<i>Tritirachium album</i> Limber	+	proteinase K	TAPROK
	<i>Tritirachium album</i>	+	proteinase R	TAPROR
	<i>Tritirachium album</i>	proT	proteinase T	TAPROT
25	<i>Aspergillus oryzae</i>	+	alkaline protease	AOALPR
	<i>Malbranchea pulchella</i>	-	thermomycin	MPTHMY
	<i>Acremonium chrysogenum</i>	alp	alkaline protease	ACALPR
	<u>Yeasts</u>			
	<i>Kluyveromyces lactis</i>	kex1	Kex1 ser. proteinase	KLKEX1
30	<i>Saccharomyces cerevisiae</i>	kex2	Kex2 ser. proteinase	SCKEX2
	<i>Saccharomyces cerevisiae</i>	prb1	protease B	SCPRB1
	<i>Yarrowia lipolytica</i>	xpr2	alk. extracell. prot.	LXPR2

SUBSTITUTE SHEET

Organism	cDNA, gene	enzyme	acronym
<u>HIGHER EUKARYOTES</u>			
<u>5 Worms</u>			
<i>Caenorhabditis elegans</i>	bli4	cuticle protease	CEBLI4
<u>Insects</u>			
<i>Drosophila</i> (fruit fly)	fur1	furin 1	DMFUR1
<i>Drosophila</i> (fruit fly)	fur2	furin 2	DMFUR2
<u>10 Plants</u>			
<i>Cucumis melo</i> (melon)	-	cucumisin	CMCUCU
<u>Mammals</u>			
Human (also rat, mouse)	fur	furin	HSFURI
Human (also mouse)	+	insulinoma PC2 prot.	HSIPC2
<u>15 Mouse</u>			
Human	+	pituitary PC3 prot.	MMPPC3
	+	tripeptidyl peptid.II	HSTPP

References used for Table I

20

References to amino acid sequences (GenBank®/EMBL Data Bank accession numbers are shown in brackets):

- ARB172 Kamekura and Seno, (1990) *Biochem. Cell Biol.* 68 352-359 (amino acid sequencing of mature protease residues 1-35; residue I4 not determined).
- 25 BSS168 Stahl. and Ferrari. (1984) *J. Bacteriol.* 158, 411-418 (K01988). Yoshimoto, Oyama et al. (I488) *J. Biochem.* 103, 1060-1065 (the mature subtilisin from *B.subtilis* var. *amylosacchariticus* differs in having T130S and T162S). Svendsen, et al. (1986) *FEBS Lett.* 196, 228-232 (PIR A23624; amino acid sequencing; the mature alkaline mesentericopeptidase From *B. mesentericus* differs in having S85A, A88S, S89A. S183A and N259S).
- 30

SUBSTITUTE SHEET

- BASBPN Wells, et al. (1983) *Nucl.Acids Res.* 11 7911-7925 (X00165). Vasantha et al., (1984) *J. Bacteriol.* 159 811-814(K02496).
- BSSDY Nedkov et al. (1983) *Hoppe-Seyler's Z. Physiol. Chem.* 364 1537-1540 (PIR A00969; amino acid sequencing).
- 5
- BLSCAR Jacobs et al. (1985) *Nucleic Acids Res.* 13 8913-8926 (X03341). Smith et al. (1968) *J. Biol. Chem.* 243 2184-2191 (PIR A00968; amino acid sequencing; mature protease sequence differs in having T103S, P129A, S158N, N161S and S212N).
- 10
- BLS147 Hastrup et al. (1989) PCT Patent Appl. WO 8906279. Pub. July 13 1989. (Esperase® from *B. lentus*). Takami et al. (1990) *Appl. Microbiol. Biotechnol.*, 33 519-523 (amino acid sequencing of mature alkaline protease residues 1-20 from *Bacillus* sp. no. AH-101; this sequence differs from BLS147 in having N11S).
- 15
- BABP92 van der Laan et al. (1991) *Appl. Environ. Microbiol.* 57 901-909. (Maxacal®). Hastrup et al. (1989) PCT Patent Appl. WO 8906279. Pub. 13 Jul 1989. (subtilisin 309. Savinase®, from *B. lentus* differs only in having N87S). Godette et al. (1991) Abstracts 5th Protein Society Symposium, June 6, Baltimore: abstract M8 (a high-alkaline protease from *B.lentus* differs in having N87S, S99D, S101R, S103A, V104I and G159S).
- 25
- BDSM48 Rettenmaier et al. (1990) PCT Patent Appl. WO 90/04022. Publ. April 19, 1990.
- 30 BYSYAB Kaneko et al. (1989) *J. Bacteriol.* 171 5232-5236 (M28537).
- BSEPR Sloma et al. (1988) *J. Bacteriol.* 170 5557-5563 (M22407). Bruckner (1990) *Mol. Gen. Genet.* 221 486-490 (X53307).

SUBSTITUTE SHEET

	BSBPF	Sloma et al. (1990) <i>J. Bacteriol.</i> 172 1470-1477 (M29035; corrected). Wu et al. (1990) <i>J. Biol.Chem.</i> 265 6845-6850 (J05400; this sequence differs in having A169V and 586 less C-terminal residues due to a frameshift).
5	BSISP1	Koide et al. (1986) <i>J. Bacteriol.</i> 167 110-116 (M13760).
	BSIA50	Strongin et al. (1978) <i>J. Bacteriol.</i> 133 1401-1411 (amino acid sequencing of mature protease residues 1-54; residues 3. 39, 40. 45, 46, 49 and 50 not determined).
10		
	BTFINI	Chestukhina et al. (1985) <i>Biokhimiya</i> 50 1724-1730 (amino acid sequencing of mature protease residues 1-14 from <i>B. thuringiensis</i> variety <i>israeliensis</i> , and residues 1-16 and 223-243 from variety <i>finitimus</i>). Kunitate et al. (1989) <i>Agric. Biol. Chem.</i> 53 3251-3256 (amino acid sequencing of mature protease residues 6-20 from variety <i>kurstaki</i> . BTKURS).
15		
	20 BCESPR	Chestukhina et al. (1985) <i>Biokhimiya</i> 50 1724-1730 (amino acid sequencing of mature residues 1-16 and 223-243).
	NDAPII	Tsujibo et al. (1990) <i>Agric. Biol. Chem.</i> 54 2177-2179 (amino acid sequencing of mature residues 1-26).
25		
	TVTHER	Meloun et al. (1985) <i>FEBS Lett.</i> 183 195-200 (PIR A00973; amino acid sequencing of mature protease residues 1-274).
	EFCYLA	Segarra et al. (1991) <i>Infect. Immun.</i> 59 1239-1246.
30	SEEP1P	Schnell et al. (1991) personal communication (Siezen et al. (<i>supra</i>)).
	SPSCPA	Chen et al. (1990) <i>J. Biol. Chem.</i> 265 3161-3167 (J05224).

SUBSTITUTE SHEET

DNEBPR	Kortt et al. (1991) Abstracts 5th Protein Society Symposium, June 22-26, Baltimore.abstract S76.
LLSK11	Vos et al. (1989) <i>J. Biol. Chem.</i> 264 13579-13585 (J04962). Kok et al. (1988) <i>Appl. Environ. Microbiol.</i> 54 231-238 (M24767; the sequence from strain Wg2 differs in 44 positions, including 18 differences in the protease domain, and a deletion of residues 1617-1676). Kiwaki et al. (1989) <i>Mol.Microbiol.</i> 3 359-369 (X14130; the sequence from strain NCD0763 differs in 46 positions, including 22 in the protease domain, and a deletion of residues 1617-1676).
5	
10	
XCEXPR	Liu et al. (1990) <i>Mol. Gen. Genet.</i> 220 433-440.
SMEXSP	Yanagida et al. (1986) <i>J. Bacteriol.</i> 166 937-994 (M13469).
15	
TAAQUA	Terada et al. (1990) <i>J. Biol. Chem.</i> 265 6576-6581 (J05414).
TRT41A	McHale et al. (1990) Abstracts 5th Eur. Congr. Biotechn. Christiansen ,Munck and Villadsen (eds), Munksgaard Int. Publishers, Copenhagen.
20	
VAPROA	Deane et al. (1989) <i>Gene</i> 76 281-288 (M25499).
SRESPD	Lavrenova et al. (1984) <i>Biochemistry USSR.</i> 49 447-454 (amino acid sequencing of residues 1-23; residues 13, 18 and 19 not determined).
25	
AVPRCA	Maldener et al (1991) <i>Mol. Gen. Genet.</i> 225 113-120 (the published sequence has 28 uncertain residues near position 200-210 due to a frameshift reading error).
TAPROK	Gunkel and Gassen (1989) <i>Eur. J. Biochem.</i> 179 185-194 (X14688/XI4689). Jany et al. (1986) <i>J. Biol. Chem.Hoppe-Seyler</i> 367 87(PIR A24541; amino acid sequencing; mature protease differs in having S745G, SILST204-208DSL and VNLL264-267FNL).
30	

SUBSTITUTE SHEET

TAPROR	Samal et al. (1990) <i>Mol. Microbiol.</i> 4 1789-1792 (X56116).
TAPROT	Samal et al. (1989) <i>Gene</i> 85 329-333.
AOALPR	Tatsumi et al. (1989) <i>Mol. Gen. Genet.</i> 219 33-38.
5	Cheevadhanarah et al. (1991) EMBL Data Library (X54726).
MPTHMY	Gaucher and Stevenson (1976) <i>Methods Enzymol.</i> 45 415-433 (amino acid sequencing of residues 1-28, and hexapeptide LSGTSM with active site serine).
10 ACALPR	Isogai et al. (1991) <i>Agric. Biol. Chem.</i> 55 471-477. Stepanov et al. (1986) <i>Int. J. Biochem.</i> 18 369-375 (amino acid sequencing of residues 1-27: the mature protease differs in having H13[1]Q, R13[2]N and S13[6]A).
15 KLKEX1	Tanguy-Rougeau, Wesolowski-Louvel and Fukuhara (1988) <i>FEBS Lett.</i> 234 464-470 (X07038).
SCKEX2	Mizuno et al. (1988) <i>Biochem. Biophys. Res. Commun.</i> 156 246-254 (M24201).
SCPRB1	Moehle et al. (1987) <i>Mol. Cell. Biol.</i> 7 4390-4399 (M18097).
20	
YLXYPR2	Davidow et al. (1987) <i>J. Bacteriol.</i> 169 4621-4629 (M17741). Matoba et al. (1988) <i>Mol. Cell Biol.</i> 8 4904-4916 (M23353).
CEBL14	Peters and Rose (1991) <i>The Worm Breeder's Gazette</i> 11 28.
25	
DMFUR1	Roebroek et al. (1991) <i>FEBS Lett.</i> 289 133-137 (X59384).
DMFUR2	Roebroek et al. (1992) 267 17208-17215.
CMCUCU	Kaneda et al. (1984) <i>J. Biochem.</i> 95 825-829 (amino acid sequencing of octapeptide NIISGTSM with active site serine).
30	
HSFURI	van den Ouweland et al. (1990) <i>Nucl. Acids Res.</i> 18 664 (X04329) (the sequence of mouse furin differs in 51 positions, including five in the catalytic

SUBSTITUTE SHEET

- domain: A15E, Y21F, S223F, A232V and N258[2]D).
 Misumi et al. (1990) *Nucl. Acids Res.* 18 6719
 (X55660: the sequence of rat furin differs in 49
 positions, including three in the catalytic domain:
 5 A15E, Y21F, H24R).
 HSIPC2 Smeekens and Steiner (1990) *J. Biol. Chem.* 265
 2997-3000 (J05252). Seidah et al. (1990) *DNA Cell
 Biol.* 9 415-424 (the sequence of mouse pituitary
 PC2 protease differs in 23 positions, including
 10 seven in the protease domain: I4F, S42[2]Y, E45D,
 N76S, D133E, V134L and G239[1]D).
 MMPPC3 Smeekens et al. (1991) *Proc. Natl. Acad. Sci. USA*
 88 340-344 (M58507). Seidah et al. (1990) *DNA Cell
 Biol.* 9 415-424 (M55668/M55669; partial sequence).
 15 HSTPP Tomkinson and Jonsson (1991) *Biochemistry* 30 168-
 174 (J05299).

BRIEF DESCRIPTION OF THE FIGURES

20

In the drawings, Fig. 1 shows an alignment of a number of the
 subtilases mentioned in Table I;

Fig. 2 is a 3-dimensional representation of subtilisin 309
 25 showing the location of the hydrophobic domain and some of the
 amino acid residues in the vicinity thereof to be substituted
 according to the invention.

30

DETAILED DESCRIPTION OF THE INVENTION

It has surprisingly been found that the storage stability
 and/or improved performance in detergents of subtilases
 generally is improved when amino acid residues situated in, or
 35 in the vicinity of a hydrophobic domain comprising the residues

SUBSTITUTE SHEET

P129, P131, I165, Y167, Y171 of subtilisin 309 are substituted for a more hydrophobic residue. The residues in question are especially E136, G159, S164, R170, A194, and G195.

5 Further, said variant exhibits a particularly high improved stability in liquid detergents and in detergents in a shaped solid form.

Fig. 2 shows the hydrophobic domain in subtilisin 309 and
10 residues in the vicinity thereof a number of which are to be substituted in order to increase the hydrophobicity of the domain. This may be achieved by substituting hydrophobic residues for non-hydrophobic residues and/or by substituting residues to become even more hydrophobic than in the parent
15 enzyme.

The same principle applies to the corresponding domain in other subtilases, the identification of which is within the skills of the average person working in this technical field. Graphic
20 representations like the one in Fig. 2 can be produced for other subtilases to determine the target residues to be substituted according to the invention.

A number hereof is indicated in Table II below:

25

SUBSTITUTE SHEET

Table II

residues in hydrophobic domain and the vicinity thereof

Pos\Enz.	BASBPN	BLSCAR	BLS309	BLS147	TVTHER
domain					
129	P	A	P	T	T
131	G	G	P	G	G
165	V	I	I	V	P
167	Y	Y	Y	Y	Y
171	Y	Y	Y	Y	Y
Vicinity					
136	K	K	E	E	Q
159	S	S	G	G	T
164	T	T	S	G	A
170	K	K	R	R	Y
194	P	A	A	P	S
195	E	E	G	E	V

5 Table II was constructed using the alignment shown in Fig. 2. It is obvious that similar or larger tables covering other subtilases may easily be produced by the skilled person.

Consequently the invention relates to subtilase variants in
 10 which the amino acid sequence has been changed through mutating the gene of the subtilisin enzyme, which it is desired to modify (the parent enzyme or gene), in the codon responsible for the expression of the amino acid residue in positions 129, 131, 165, 167, 171, 136, 159, 164, 170, 194, and 195, which
 15 residues are more hydrophobic than the residue(s) in the parent

SUBSTITUTE SHEET

enzyme, especially such hydrophobic residues that comprise a relatively long hydrophobic side chain, such as Ile, Leu, and Val, whereby, when the mutated gene is expressed, the amino acid residue is substituted by a more hydrophobic residue, which increases the hydrophobicity of the domain as such.

Hydrophobic amino acid residues are generally the following:

Val (V), Ile (I), Leu (L), Met (M), Phe (F), Pro (P) and Trp(W). Among these Val, Ile and Leu are preferred.

10

By looking at Table II and applying the principle of the invention a number of candidates for substitution becomes clear.

15 For both BASBPN and BLSCAR it seems appropriate to make substitutions in positions 129, 131, 136, 159, 164, 167, 170, 171 and 195. In BLS309 positions 136, 164, 167, and 170, 171 would be the first choices, and positions 159 and 195 also would be a second choice. In BLS147 positions 129, 131, 136, 20 167, 170, 171 and 195 are the first choice, while positions 159 and 164 are second. Finally, in TVTHER positions 129, 131, 136, 167, 171 and 194 are the first choices, with 164 as a second one.

25 According to the invention it would entail an advantage to substitute the Gly residues in the hydrophobic domain to bulkier and more hydrophobic residues.

Such considerations apply for any hydrophilic or hydrophobic 30 residue that may occupy any of the above mentioned position, meaning that any increase in hydrophobicity seems to be advantageous. This means that e.g. a very hydrophilic residue such as the charged residues Arg (R), Asp (D), Glu (E) or Lys (K) may be substituted by any residue that is less hydrophilic. 35 Such less hydrophilic residues comprises the residues Gly (G),

Cys (C), Ser (S), Ala (A), Thr (T), Tyr (Y), Gln (Q), His (H) or Asn (N). It also means that a Tyr(Y) may be substituted by a more hydrophobic residue such as Phe(F), Leu(L), or Ile(I).

5 Similar considerations can be applied to other subtilases having a hydrophobic domain in this part of the surface of the enzyme.

In the context of this invention a subtilase is defined in
10 accordance with Siezen et al. *supra*. In a more narrow sense, applicable to many embodiments of the invention, the subtilases of interest are those belonging to the subgroups I-S1 and I-S2. In a more specific sense, many of the embodiments of the invention relate to serine proteases of gram-positive bacteria
15 which can be brought into substantially unambiguous homology in their primary structure, with the subtilases listed in Table I above.

The present invention also comprises any one or more
20 substitutions in the above mentioned positions in combination with any other substitution, deletion or addition to the amino acid sequence of the parent enzyme. Especially combinations with other substitutions known to provide improved properties to the enzyme are envisaged.

25

Such combinations comprise the positions: 222 (improve oxidation stability), 218 (improves thermal stability), substitutions in the Ca-binding sites stabilising the enzyme, e.g. position 76, and many other apparent from the prior art.

30

Furthermore combinations with the variants mentioned in EP 405 901 are also contemplated specifically.

SUBSTITUTE SHEET

VARIANTS

A: Single variants:

Subtilisin BPN', Subtilisin Carlsberg, Subtilisin 168, and Subtilisin DY variants:

- 5 A129V, A129I, A129L, A129M, A129F
 G131V, G131I, G131L, G131M, G131F
 K136V, K136I, K136L, K136M, K136F,
 S159V, S159I, S159L, S159M, S159F,
 T164V, T164I, T164L, T164M, T164F,
- 10 Y167V, Y167I, Y167L, Y167M, Y167F
 K170V, K170I, K170L, K170M, K170F,
 Y171V, Y171I, Y171L, Y171M, Y171F
 A194V, A194I, A194L, A194M, A194F
 E195V, E195I, E195L, E195M, E195F,

15

Thermitase variants:

- A129V, A129I, A129L, A129M, A129F
 G131V, G131I, G131L, G131M, G131F
 Q136V, Q136I, Q136L, Q136M, Q136F,
- 20 T159V, T159I, T159L, T159M, T159F,
 A164V, A164I, A164L, A164M, A164F,
 Y167V, Y167I, Y167L, Y167M, Y167F
 Y171V, Y171I, Y171L, Y171M, Y171F
 Y170V, Y170I, Y170L, Y170M, Y170F,
- 25 S194V, S194I, S194L, S194M, S194F,

Subtilisin 309, Subtilisin 147, and Bacillus PB92 protease variants:

- T129V, T129I, T129L, T129M, T129F
- 30 G131V, G131I, G131L, G131M, G131F
 E136V, E136I, E136L, E136M, E136F,
 G159V, G159I, G159L, G159M, G159F,
 G164V, G164I, G164L, G164M, G164F, (BLS147)
 S164V, S164I, S164L, S164M, S164F, (BLS309 AND BAPB92)
- 35 Y167A, Y167H, Y167N, Y167P, Y167C, Y167W,

SUBSTITUTE SHEET

Y167Q, Y167S, Y167T, Y167G, Y167V, Y167I, Y167L, Y167M, Y167F
 R170W, R170A, R170H, R170N, R170P, R170Q, R170S, R170T, R170Y
 (disclaimed for BLS309), R170V (disclaimed for BAPB92), R170I
 (disclaimed for BAPB92), R170L, R170M (disclaimed for
 5 BAPB92), R170F, R170G, R170C,

Y171A, Y171H, Y171N, Y171P, Y171C, Y171W, Y171Q, Y171S,
 Y171T, Y171G, Y171V, Y171I, Y171L, Y171M, Y171F

A194V, A194I, A194L, A194M, A194F, (BLS309 AND BAPB92)

P194V, P194I, P194L, P194M, P194F, (BLS147)

10 E195V, E195I, E195L, E195M, E195F, (BLS147)

G195V, G195I, G195L, G195M, G195F, (BLS309 AND BAPB92)

B: Combination Variants:

15 Any of the above variants are contemplated to prove
 advantageous if combined with other variants in any of the
 positions:

27, 36, 57, 76, 97, 101, 104, 120, 123, 206, 218, 222, 224, 235
 and 274.

20

Specifically the following BLS309 and BAPB92 variants are
 considered appropriate for combination:

K27R, *36D, S57P, N76D, G97N, S101G, V104A, V104N, V104Y,
 H120D, N123S, A194P, Q206E, N218S, M222S, M222A, T224S, K235L

25 and T274A.

Also such variants comprising any one or two of the
 substitutions X167V, X167M, X167F, X167L, X167I, X170V, X170M,
 X170F, X170L, and/or X170I in combination with any one or more

30 of the other substitutions, deletions and/or insertions
 mentioned above are advantageous.

Furthermore variants comprising any of the variants
 V104N+S101G, K27R+V104Y+N123S+T274A, or N76D+V104A or other

35 combinations of these mutations (V104N, S101G, K27R, V104Y,

SUBSTITUTE SHEET

N123S, T274A, N76D, V104A), in combination with any one or more of the substitutions, deletions and/or insertions mentioned above are deemed to exhibit improved properties.

5 Specific combinations to be mentioned are:

- a) S57P+R170L
- a') S57P+R170I
- b) R170L+N218S
- b') R170I+N218S
- 10 c) S57P+R170L+N218S
- c') S57P+R170I+N218S
- c") S57P+V104Y+R170L+N218S
- c'") S57P+V104Y+R170I+N218S
- d) R170L+N218S+M222A
- 15 d') R170I+N218S+M222S
- d") R170L+N218S+M222A
- d'") R170I+N218S+M222S
- e) S57P+R170L+S188P+A194P
- e') S57P+R170I+S188P+A194P
- 20 f) Y167L+R170L
- f') Y167L+R170I
- g) Y167I+R170L
- g') Y167I+R170I
- h) N76D+R170L+N218S
- 25 h') N76D+R170I+N218S
- i) S57P+N76D+R170L+N218S
- i') S57P+N76D+R170I+N218S
- j) N76D+R170L+N218S+M222A
- j') N76D+R170I+N218S+M222S
- 30 j") N76D+R170L+N218S+M222A
- j'") N76D+R170L+N218S+M222S
- k) S57P+R170I+S188P+A194P+N218S
- k') S57P+R170I+S188P+A194P+N218S
- l) *36D+N76D+H120D+R170L+G195E+K235L
- 35 l') *36D+N76D+H120D+R170I+G195E+K235L

SUBSTITUTE SHEET

1") *36D+N76D+H120D+Y167I+R170L+G195E+K235L
 1'') *36D+N76D+H120D+Y167I+R170I+G195E+K235L
 m) N76D+H120D+R170L+G195E+K235L
 m') N76D+H120D+R170I+G195E+K235L
 5 m'') N76D+H120D+Y167I+R170L+G195E+K235L
 m''') N76D+H120D+Y167I+R170I+G195E+K235L
 n) *36D+G97N+V104Y+H120D+R170L+A194P+G195E+K235L
 n') *36D+G97N+V104Y+H120D+R170I+A194P+G195E+K235L
 o) S57P+R170L+Q206E
 10 o') S57P+R170I+Q206E
 p) R170L+Q206E
 p') R170I+Q206E
 q) Y167I+R170L+Q206E
 q') Y167I+R170I+Q206E
 15 r) Y167F+R170L
 r') Y167F+R170I
 t) Y167I+R170L+A194P
 t') Y167I+R170I+A194P
 t'') Y167L+R170L+A194P
 20 t''') Y167L+R170I+A194P
 u) Y167I+R170L+N218S
 u') Y167I+R170I+N218S
 u'') Y167L+R170L+N218S
 u''') Y167L+R170I+N218S
 25 v) Y167I+R170L+A194P+N218S
 v') Y167I+R170I+A194P+N218S
 v'') Y167L+R170L+A194P+N218S
 v''') Y167L+R170I+A194P+N218S
 x) R170L+P131V
 30 x') R170I+P131V
 y) *36D+Y167I+R170L
 y') *36D+Y167I+R170I
 z) Y167I+Y171I
 aa) Y167V+R170L
 35 aa') Y167V+R170I

SUBSTITUTE SHEET

bb) R170L+Y171I
 bb') R170I+Y171L
 bb'') R170L+Y171L
 bb''') R170I+Y171I
 5 cc) Y167I+Y171L+N218S
 cc') Y167I+Y171I+N218S

DETERGENT COMPOSITIONS COMPRISING THE MUTANT ENZYMES

10 The present invention also comprises the use of the mutant enzymes of the invention in cleaning and detergent compositions and such compositions comprising the mutant subtilisin enzymes. Such cleaning and detergent compositions can in principle have any physical form, but the subtilase variants are preferably
 15 incorporated in liquid detergent compositions or in detergent compositions in the form of bars, tablets, sticks and the like for direct application, wherein they exhibit improved enzyme stability or performance.

20 Among the liquid compositions of the present invention are aqueous liquid detergents having for example a homogeneous physical character, e.g. they can consist of a micellar solution of surfactants in a continuous aqueous phase, so-called isotropic liquids.

25

Alternatively, they can have a heterogeneous physical phase and they can be structured, for example they can consist of a dispersion of lamellar droplets in a continuous aqueous phase, for example comprising a deflocculating polymer having a
 30 hydrophilic backbone and at least one hydrophobic side chain, as described in EP-A-346 995 (Unilever) (incorporated herein by reference). These latter liquids are heterogeneous and may contain suspended solid particles such as particles of builder materials e.g. of the kinds mentioned below.

35

SUBSTITUTE SHEET

Concerning powder detergent compositions such compositions comprise in addition to any one or more of the subtilisin enzyme variants in accordance to any of the preceding aspects of the invention alone or in combination any of the usual 5 components included in such compositions which are well-known to the person skilled in the art.

Such components comprise builders, such as phosphate or zeolite builders, surfactants, such as anionic, cationic, non-ionic or 10 zwitterionic type surfactants, polymers, such as acrylic or equivalent polymers, bleach systems, such as perborate- or amino-containing bleach precursors or activators, structurants, such as silicate structurants, alkali or acid to adjust pH, humectants, and/or neutral inorganic salts.

15

Furthermore, a number of other ingredients are normally present in the compositions of the invention, such as:

- A. Cosurfactants
- 20 B. Tartrate Succinate Builder
- C. Neutralization System
- D. Suds Suppressor
- E. Other Enzymes
- F. Other Optional Components

25

The weight ratio of anionic surfactant to nonionic surfactant is preferably from 1:1 to 5:1. The compositions have a pH in a 10% by weight solution in water at 20°C of from 7.0 to 9.0, a Critical Micelle Concentration of less than or equal to 200 30 ppm, and an air/water Interfacial Tension at the Critical Micelle Concentration of less than or equal to 32 dynes/cm at 35°C in distilled water. The compositions are preferably clear, homogeneous and phase stable, and have good cleaning performance and enzyme stability.

35

SUBSTITUTE SHEET

VARIOUS COMPONENTS:

1. Anionic Surfactant

5 The compositions of the present invention contain from about 10% to about 50%, preferably from about 15% to about 50%, more preferably from about 20% to 40%, and most preferably from 20% to about 30%, by weight of a natural or synthetic anionic surfactant. Suitable natural or synthetic anionic surfactants
10 are e.g. soaps and such as disclosed in U.S. Patent 4,285,841, and in U.S. Patent 3,929,678.

Useful anionic surfactants include the water-soluble salts, particularly the alkali metal, ammonium and alkylolammonium
15 (e.g., monoethanolammonium or triethanolammonium) salts, of organic sulfuric reaction products having in their molecular structure an alkyl group containing from about 10 to about 20 carbon atoms and a sulfonic acid or sulfuric acid ester group. (Included in the term "alkyl" is the alkyl portion of aryl
20 groups.) Examples of this group of synthetic surfactants are the alkyl sulfates, especially those obtained by sulfating the higher alcohols (C_8 - C_{16} carbon atoms) such as those produced by reducing the glycerides of tallow or coconut oil; and the alkylbenzene sulfonates in which the alkyl group contains from
25 about 9 to about 15 carbon atoms, in straight chain or branched chain configuration, e.g., those of the type described in U. S. Patents 2,220,099 and 2,477,383. Especially valuable are linear straight chain alkylbenzene sulfonates in which the average number of carbon atoms in the alkyl group is from about 11 to
30 14.

Other anionic surfactants herein are the water-soluble salts of: paraffin sulfonates containing from 8 to about 24 (preferably about 12 to 18) carbon atoms; alkyl glyceryl ether sulfonates, especially those ethers of C_8 - C_{18} alcohols (e.g., those
35

SUBSTITUTE SHEET

derived from tallow and coconut oil); alkyl phenol ethylene oxide ether sulfates containing from 1 to about 4 units of ethylene oxide per molecule and from 8 to 12 carbon atoms in the alkyl group; and alkyl ethylene oxide ether sulfates 5 containing 1 to 4 units of ethylene oxide per molecule and from 10 to 20 carbon atoms in the alkyl group.

Other useful anionic surfactants include the water-soluble salts of esters of a-sulfonated fatty acids containing from 6 10 to 20 carbon atoms in the fatty acid group and from 1 to 10 carbon atoms in the ester group; water-soluble salts of 2-acyloxy-alkane-1-sulfonic acids containing from 2 to 9 carbon atoms in the acyl group and from 9 to 23 carbon atoms in the alkane moiety; water-soluble salts of olefin sulfonates 15 containing from 12 to 24 carbon atoms; and b-alkyloxy alkane sulfonates containing from 1 to 3 carbon atoms in the alkyl group and from 8 to 20 carbon atoms in the alkane moiety.

Preferred anionic surfactants are soaps, the C₁₀-C₁₈ alkyl 20 sulfates and alkyl ethoxy sulfates containing an average of up to 4 ethylene oxide units per mole of alkyl sulfate, C₁₁-C₁₃ linear alkyl benzene sulfonates, and mixtures thereof.

2. Nonionic Surfactant

25

Another optional ingredient is from 2% to 14% preferably from 2% to 8%, most preferably from 3% to 5% by weight, of an optionally ethoxylated nonionic surfactant. The weight ratio of natural or synthetic anionic surfactant (on an acid basis) to 30 nonionic surfactant is from 1:1 to 5:1 preferably from 2:1 to 5:1, most preferably from 3:1 to 4:1. This is to ensure the formation and adsorption of sufficient hardness surfactants at the air/water interface to provide good greasy/oily soil removal.

35

SUBSTITUTE SHEET

The optionally ethoxylated nonionic surfactant is of the formula $R^1(OC_2H_4)_n OH$, wherein R^1 is a C_{10} - C_{16} alkyl group or a C_8 - C_{12} alkyl phenyl group, n is from 3 to 9, and said nonionic surfactant has an HLB (Hydrophilic-Lipophilic Balance) of from 5 6 to 14, preferably from 10 to 13. These surfactants are more fully described in U.S. Patents 4,285,841, and 4,284,532. Particularly preferred are condensation products of C_{12} - C_{15} alcohols with from 3 to 8 moles of ethylene oxide per mole of alcohol, e.g., C_{12} - C_{13} alcohol condensed with about 6.5 moles of 10 ethylene oxide per mole of alcohol. Other nonionic surfactants to be mentioned are APG, EGE, and glucamide surfactants.

3. Detergency Builder

15 Among the usual detergent ingredients which may be present in usual amounts in the detergent compositions of this invention are the following: The compositions may be built or unbuilt and may be of the zero-P type (i.e. not containing any phosphorus containing builders). Thus, the composition may contain 20 in the aggregate for example from 1-50%, e.g. at least about 5% and often up to about 35-40% by weight, of one or more organic and/or inorganic builders. Typical examples of builders include those already mentioned above, and more broadly include alkali metal ortho, pyro, and tripolyphosphates, alkali metal carbon- 25 ates, either alone or in admixture with calcite, alkali metal citrates, alkali metal nitrilotriacetates, carboxymethyloxysuccinates, zeolites, polyacetalcarboxylates, and so on.

More specifically the compositions herein contain from 5% to 30 20%, preferably from 10% to 15%, by weight of a detergency builder which can be a fatty acid containing from 10 to 18 carbon atoms and/or a polycarboxylate, zeolite, polyphosphonate and/or polyphosphate a builder. Preferred are from 0 to 10% (more preferably from 3% to 10%) by weight of saturated fatty 35 acids containing from 12 to 14 carbon atoms, along with from 0

SUBSTITUTE SHEET

to 10%, more preferably from 2% to 8%, most preferably from 2% to 5%, by weight of a polycarboxylate builder, most preferably citric acid, in a weight ratio of from 1:1 to 3:1.

5 Since the proteolytic enzymes herein appear to provide optimum storage stability benefits versus other enzymes when the builder to water hardness ratio is close to one, the compositions preferably contain sufficient builder to sequester from 2 to 10, preferably from 3 to 8, grains per gallon of
10 hardness.

Suitable saturated fatty acids can be obtained from natural sources such as plant or animal esters (e.g., palm kernel oil, palm oil and coconut oil) or synthetically prepared (e.g., via
15 the oxidation of petroleum or by hydrogenation of carbon monoxide via the Fisher-Tropsch process). Examples of suitable saturated fatty acids for use in the compositions of this invention include capric, lauric, myristic, coconut and palm kernel fatty acid. Preferred are saturated coconut fatty acids;
20 from 5:1 to 1:1 (preferably about 3:1) weight ratio mixtures of lauric and myristic acid; mixtures of the above with minor amounts (e.g., 1%-30% of total fatty acid) of oleic acid; and palm kernel fatty acid.

25 The compositions herein preferably also contain the polycarboxylate, polyphosphonate and polyphosphate builders described in U.S. Patent 4,284,532, Water-soluble polycarboxylate builders, particularly citrates, are preferred of this group. Suitable polycarboxylate builders include the
30 various aminopolycarboxylates, cycloalkane polycarboxylates, ether polycarboxylates, alkyl polycarboxylates, epoxy polycarboxylates, tetrahydrofuran polycarboxylates, benzene polycarboxylates, and polyacetal polycarboxylates.

SUBSTITUTE SHEET

Examples of such polycarboxylate builders are sodium and potassium ethylenediaminetetraacetate; sodium and potassium nitrilotriacetate; the water-soluble salts of phytic acid, e.g., sodium and potassium phytates, disclosed in U.S. Patent 5 1,739,942, the polycarboxylate materials described in U.S. Patent 3,364,103; and the water-soluble salts of polycarboxylate polymers and copolymers described in U.S. Patent 3,308,067.

10 Other useful detergency builders include the water-soluble salts of polymeric aliphatic polycarboxylic acids having the following structural and physical characteristics: (a) a minimum molecular weight of about 350 calculated as to the acid form; (b) an equivalent weight of 50 to 80 calculated as to
15 acid form; (3) at least 45 mole percent of the monomeric species having at least two carboxyl radicals separated from each other by not more than two carbon atoms: (d) the site of attachment of the polymer chain of any carboxyl-containing radical being separated by not more than three carbon atoms
20 along the polymer chain from the site of attachment of the next carboxyl-containing radical. Specific examples of such builders are the polymers and copolymers of itaconic acid, aconitic acid, maleic acid, mesaconic acid, fumaric acid, methylene malonic acid, and citraconic acid.

25

Other suitable polycarboxylate builders include the water-soluble salts, especially the sodium and potassium salts, of mellitic acid, citric acid, pyromellitic acid, benzene pentacarboxylic acid, oxydiacetic acid, carboxymethyloxy-
30 succinic acid, carboxymethyloxymalonic acid, cis-cyclohexane-hexacarboxylic acid, cis-cyclopentanetetracarboxylic acid and oxydisuccinic acid.

Other polycarboxylates are the polyacetal carboxylates
35 described in U.S. Patent 4,144,226, and U.S. Patent 4,146,495.

SUBSTITUTE SHEET

Other detergency builders include the zeolites, such as the aluminosilicate ion exchange material described in U.S. Patent 4,405,483.

5

Other preferred builders are those of the general formula $R-CH(COOH)CH_2(COOH)$, i.e. derivatives of succinic acid, wherein R is $C_{10}-C_{20}$ alkyl or alkenyl, preferably $C_{12}-C_{16}$, or wherein R may be substituted with hydroxyl, sulfo, sulfoxy or sulfone substituents. These succinate builders are preferably used in the form of their water soluble salts, including the sodium, potassium and alkanolammonium salts. Specific examples of succinate builders include: lauryl succinate, myristyl succinate, palmityl succinate, 2-dodecenyl succinate, and the like.

4. Proteolytic Enzyme

The enzymes of the invention can be used in well-known standard amounts in detergent compositions. The amounts may range very widely, e.g. about 0.0002-0.1, e.g. about 0.005-0.05, Anson units per gram of the detergent composition. Expressed in alternative units, the protease can be included in the compositions in amounts in the order of from about 0.1 to 100 GU/mg (e.g. 1-50, especially 5-20 GU/mg) of the detergent formulation, or any amount in a wide range centering at about 0.01-4, e.g. 0.1-0.4 KNPU per g of detergent formulation.

It may for example be suitable to use the present enzymes at the rate of about 0.25 mg of enzyme protein per litre of wash liquor, corresponding to an enzyme activity of the order of 0.08 KNPU per litre. Corresponding detergent formulations can contain the enzymes in for example an amount of the order of 0.1-0.4 KNPU/g.

35

SUBSTITUTE SHEET

Expressed differently the compositions of the present invention contain from about 0.01% to about 5%, preferably from about 0.1% to about 2%, by weight of the proteolytic enzymes of the invention.

5

The described proteolytic enzyme is preferably included in an amount sufficient to provide an activity of from 0.05 to about 1.0, more preferably from about 0.1 to 0.75, most preferably from about 0.125 to about 0.5,mg of active enzyme per gram of
10 composition.

The enzyme component may be added to the other components in any convenient form, such as in the form of a solution, slurry, LDP slurry, or crystals.

15

5. Enzyme Stabilization System

The liquid detergents according to the present invention may comprise An enzyme stabilization system, comprising calcium
20 ion, boric acid, propylene glycol and/or short chain carboxylic acids. The enzyme stabilization system comprises from about 0.5% to about 15% by weight of the composition.

The composition preferably contains from about 0.01 to about
25 50, preferably from about 0.1 to about 30, more preferably from about 1 to 20 millimoles of calcium ion per liter. The level of calcium ion should be selected so that there is always some minimum level available for the enzyme, after allowing for complexation with builders etc. in the composition. Any water-
30 soluble calcium salt can be used as the source of calcium ion, including calcium chloride, calcium formate, and calcium acetate. A small amount of calcium ion, generally from about 0.05 to 0.4 millimoles per liter, is often also present in the composition due to calcium in the enzyme slurry and formula

SUBSTITUTE SHEET

water. From about 0.03% to about 0.6% of calcium formate is preferred.

A second preferred enzyme stabilizer is polyols containing only 5 carbon, hydrogen and oxygen atoms. They preferably contain from 2 to 6 carbon atoms and from 2 to 6 hydroxy groups. Examples include propylene glycol (especially 1,2-propanediol, which is preferred), ethylene glycol, glycerol, sorbitol, mannitol, and glucose. The polyol generally represents from about 0.5% to 10 15%, preferably from about 1.5% to about 8%, by weight of the composition. Preferably, the weight ratio of polyol to any boric acid added is at least 1, more preferably at least 1.3.

The compositions preferably also contain the water-soluble, 15 short chain carboxylates described in U.S. Patent 4,318,818. The formates are preferred and can be used at levels of from about 0.05% to about 5%, preferably from about 0.2% to about 2%, most preferably from 0.4% to 1.5%, by weight of the composition. Sodium formate is preferred.

20

The compositions herein also optionally contain from about 0.25% to about 5%, most preferably from about 0.5% to about 3%, by weight of boric acid. The boric acid may be, but is preferably not, formed by a compound capable of forming boric 25 acid in the composition. Boric acid is preferred, although other compounds such as boric oxide, borax and other alkali metal borates (e.g., sodium ortho-, meta- and pyroborate, and sodium pentaborate) are suitable. [Substituted boric acids (e.g., phenylboronic acid, butane boronic acid, and p-bromo 30 phenylboronic acid) can also be used in place of boric acid.

6. Water

The liquid compositions of the present invention may be aqueous 35 liquids or non-aqueous liquids. When the are aqueous liquids,


SUBSTITUTE SHEET

they contain from about 15% to about 60%, preferably from about 25% to about 45%, by weight of water.

5 FURTHER OPTIONAL COMPONENTS

A. Cosurfactants

Optional cosurfactants for use with the above nonionic
10 surfactants include amides of the formula



15

wherein R^1 is an alkyl, hydroxyalkyl or alkenyl radical containing from 8 to 20 carbon atoms, and R^2 and R^3 are selected from the group consisting of hydrogen, methyl, ethyl, propyl, isopropyl, 2-hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl,
20 and said radicals additionally containing up to 5 ethylene oxide units, provided at least one of R^2 and R^3 contains a hydroxyl group.

Preferred amides are the C_8 - C_{20} fatty acid alkylol amides in
25 which each alkylol group contains from 1 to 3 carbon atoms, and additionally can contain up to 2 ethylene oxide units. Particularly preferred are the C_{12} - C_{16} fatty acid monoethanol and diethanol amides.

30 If used, amides are preferably present at a level such that the above ethoxylated nonionic surfactant and amide surfactant is in a weight ratio of from 4:1 to 1:4, preferably from 3:1 to 1:3.


SUBSTITUTE SHEET

Preferred and optional cosurfactants, used at a level of from 0.15% to 1%, are the quaternary ammonium, amine and amine oxide surfactants described in U.S. Patent 4,507,219.


5 Of the above, the C₁₀-C₁₄ alkyl trimethylammonium salts are preferred, e.g., decyl trimethylammonium methylsulfate, lauryl trimethylammonium chloride, myristyl trimethylammonium bromide, and coconut trimethylammonium chloride and methylsulfate. From 0.2% to 0.8% of monoalkyl trimethylammonium chloride is
10 preferred.

B. Tartrate Succinate Builder

The compositions herein preferably contain from 0 to about 10%,
15 preferably from 0 to about 6%, by weight on an acid basis, of a tartrate succinate builder material selected from the group consisting of:



wherein X is a salt-forming cation;



wherein X is a salt-forming cation; and

30 iii) mixtures thereof.

The tartrate succinate compounds used herein are described in U.S. Patent 4,663,071.

35

SUBSTITUTE SHEET

C. Neutralization System

The present compositions can also optionally contain from about 0 to about 0.04 moles, preferably from about 0.01 to 0.035 moles, more preferably from about 0.015 to about 0.03 moles, per 100 grams of composition of an alkanolamine selected from the group consisting of monoethanolamine, diethanolamine, triethanolamine, and mixtures thereof. Low levels of the alkanolamines, particularly monoethanolamine, are preferred to enhance product stability, detergency performance, and odour. However, the amount of alkanolamine should be minimized for best chlorine bleach compatibility.

In addition, the compositions contain sodium ions, and preferably potassium ions, at a level sufficient to neutralize the anionic species and provide the desired product pH.

D. Suds Suppressor

Another optional component for use in the liquid detergents herein is from 0 to about 1.5%, preferably from about 0.5% to about 1.0%, by weight of silicone based suds suppressor agent. Silicones are widely known and taught for use as highly effective suds controlling agents. For example, U.S. Patent 3,455,839 relates to compositions and processes for defoaming aqueous solutions by incorporating therein small amounts of polydimethylsiloxane fluids.

Useful suds controlling silicones are mixtures of silicone and silanated silica as described, for instance, in German Patent Application DOS 2,124,526.

Silicone defoamers and suds controlling agents have been successfully incorporated into granular detergent compositions

SUBSTITUTE SHEET

by protecting them from detergent surfactants as in U.S. Patent 3,933,672, and in U.S. Patent 4,652,392.

A preferred silicone based suds suppressor for use herein is a suds suppressing amount of a suds controlling agent consisting essentially of:

- (i) polydimethylsiloxane fluid having a viscosity of from about 20 cs. to about 1500 cs. at 25°C;
- 10 (ii) from about 5 to about 50 parts per 100 parts by weight of
 - (i) of siloxane resin composed of $(\text{CH}_3)_3\text{SiO}_{1/2}$ units and SiO_2 units in a ratio of from $(\text{CH}_3)_3\text{SiO}_{1/2}$ units and to SiO_2 units of from about 0.6:1 to about 1.2:1; and
- 15 (iii) from about 1 to about 20 parts per 100 parts by weight of (i) of a solid silica gel.

By "suds suppressing amount" is meant that the formulator of the composition can select an amount of this suds controlling agent that will control the suds to the extent desired. The amount of suds control will vary with the detergent surfactant selected. For example, with high sudsing surfactants, relatively more of the suds controlling agent is used to achieve the desired suds control than with low foaming surfactants.

E. Other Enzymes

30 The detergent compositions of the invention may also contain further enzymes.

For example, lipase can usefully be added in the form of a solution or a slurry of lipolytic enzyme with carrier material

SUBSTITUTE SHEET

(e.g. as in EP Patent Publication No. 258,068 (Novo Nordisk A/S)

The added amount of lipase can be chosen within wide limits, 5 for example 50 to 30,000 LU/g per gram of the surfactant system or of the detergent composition, e.g. often at least 100 LU/g, very usefully at least 500 LU/g, sometimes preferably above 1000, above 2000 LU/g or above 4000 LU/g or more, thus very often within the range of 50-4000 LU/g, and possibly within the 10 range of 200-1000 LU/g. In this specification, lipase units are defined as they are in EP Patent Publication No. 258,068.

The lipolytic enzyme can be chosen among a wide range of lipases. In particular, the lipases described in for example 15 the following patent specifications: EP Patent Publications Nos. 214,761 (Novo Nordisk A/S), 258,068, and especially lipases showing immunological cross reactivity with antisera raised against lipase from *Thermomyces lanuginosus* ATCC 22070, EP Patent Publications Nos. 205,208 and 206,390, and especially 20 lipases showing immunological cross-reactivity with antisera raised against lipase from *Chromobacter viscosum* var lipolyticum NRRL B-3673, or against lipase from *Alcaligenes* PL-679, ATCC 31371 and FERM-P 3783, also the lipases described in specifications WO 87/00859 (Gist-Brocades) and EP Patent 25 Publication No. 204,284 (Sapporo Breweries). Suitable, in particular, are for example the following commercially available lipase preparations: Lipolase® Novo Nordisk A/S, Amano lipases CE, P, B, AP, M-AP, AML, and CES, and Meito lipases MY-30, OF, and PL, also Esterase® MM (Novo Nordisk 30 A/S), Lipozym, SP225, SP285, (all Novo Nordisk A/S) Saiken lipase, Enzeco lipase, Toyo Jozo lipase and Diosynth lipase (Trade Marks), Lumafast® (Genencor Inc.), Lipomax® (Gist-Brocades N.V.), and lipases as described in WO 94/03578 (Unilever).

35

SUBSTITUTE SHEET

Amylase can for example be used when desired, in an amount in the range of about 1 to about 100 MU (maltose units) per gram of detergent composition (or 0.014-1.4, e.g. 0.07-0.7, KNU/g (Novo units)). Amylases suitable are for example Termamyl®, and
 5 BAN (Novo Nordisk A/S). Cellulase can for example be used when desired, in an amount in the range of about 0.3 to about 35 CEVU units per gram of the detergent composition. Suitable cellulases are for example Celluzyme®, and Carezyme® (NOVO NORDISK A/S).

10

Other enzymes contemplated to be used in the present invention are oxidases and peroxidases

F. Other Optional Components

15 Other optional components for use in the liquid detergents herein include soil removal agents, soil release polymers, antiredeposition agents such as tetraethylene pentamine ethoxylate (from about 0.5% to 3%, preferably from about 1% to about 3%, by weight), suds regulants, poly vinyl pyrrolidone,
 20 carboxy methyl cellulose, clays, and hydrotropes such as sodium cumene sulfonate, opacifiers, antioxidants, bactericides, dyes, perfumes, and brighteners known in the art. Such optional components generally represent less than about 15%, preferably from about 0.5% to 10%, more preferably from about 1% to about
 25 10%, by weight of the composition.

The compositions may contain from 0% to about 8%, preferably from 0% to about 5%, by weight of a C_{12} - C_{14} alkenyl succinic acid or salt thereof. These materials are of the general formula R-
 30 $CH(COOX)CH_2(COOX)$, wherein R is a C_{12} - C_{14} alkenyl group and each X is H or a suitable cation, such as sodium, potassium, ammonium or alkanolammonium (e.g., mono-, di-, or tri-ethanolammonium).

SUBSTITUTE SHEET

Specific examples are 2-dodecenyl succinate (preferred) and 2-tetradecenyl succinate.

The compositions herein optionally contain from about 0.1% to 5 about 1%, preferably from about 0.2% to about 0.6%, by weight of water-soluble salts of ethylenediamine tetramethylenephosphonic acid, diethylenetriamine pentamethylenephosphonic acid, ethylenediamine tetraacetic acid (preferred), or diethylenetriamine pentaacetic acid (most preferred) to enhance cleaning 10 performance when pretreating fabrics.

Furthermore, the detergent compositions may contain from 1-35% of a bleaching agent or a bleach precursor or a system comprising bleaching agent and/or precursor with activator therefor.

15

Further optional ingredients are lather boosters, anti-corrosion agents, soil-suspending agents, sequestering agents, anti-soil redeposition agents, and so on.

20 The compositions herein preferably contain up to about 10% of ethanol.

G. Other Properties

25 The instant composition usually has a pH, in a 10% by weight solution in water at 20°C, of from about 7.0 to 9.0, preferably from about 8.0 to about 8.5.

The instant compositions can also have a Critical Micelle 30 Concentration (CMC) of less than or equal to 200 parts per million (ppm), and an air/water Interfacial Tension above the CMC of less than or equal to 32, preferably less than or equal to about 30, dynes per centimetre at 35°C in distilled water. These measurements are described in "Measurement of Interfacial 35 Tension and Surface Tension - General Review for Practical Man"

SUBSTITUTE SHEET

C. Weser, *GIT Fachzeitschrift für das Laboratorium*, 24 (1980) 642-648 and 734-742, FIT Verlag Ernst Giebeler, Darmstadt, and "Interfacial Phenomena - Equilibrium and Dynamic Effects", C. A. Miller and P. Neogi, Chapter 1, pp. 29-36 (1985), Marcel Dekker, Inc. New York.

The compositions of the invention can be used for the washing of textile materials, especially, but without limitation cotton and polyester based textiles and mixtures thereof. For example washing processes carried out at temperatures of about 60-65°C or lower, e.g. about 30-35°C or lower, are particularly suitable. It can be very suitable to use the compositions at a rate sufficient to provide about e.g. 0.4-0.8 g/l of surfactant in the wash liquor, although it is of course possible to use lower or higher concentrations, if desired. Without limitation it can for example be stated that a use-rate from about 1 to 10 g/l, e.g. from about 3-6 g/l, of the detergent formulation is suitable for use in the case when the formulations are substantially as in the Examples.

20

In this aspect the invention is especially related to:

a) A detergent composition formulated as an aqueous detergent liquid comprising anionic surfactant, nonionic surfactant, humectant, organic acid, caustic alkali, with a pH adjusted to a value between 9 and 10.

b) A detergent composition formulated as a non-aqueous detergent liquid comprising a liquid nonionic surfactant consisting essentially of linear alkoxylated primary alcohol, triacetin, sodium triphosphate, caustic alkali, perborate monohydrate bleach precursor, and tertiary amine bleach activator, with a pH adjusted to a value between about 9 and 10.

35

SUBSTITUTE SHEET

- c) An enzymatic liquid detergent composition formulated to give a wash liquor pH of 9 or less when used at a rate corresponding to 0.4-0.8 g/l surfactant.
- 5 d) An enzymatic liquid detergent composition formulated to give a wash liquor pH of 8.5 or more when used at a rate corresponding to 0.4-0.8 g/l surfactant.
- e) An enzymatic liquid detergent composition formulated to
10 give a wash liquor ionic strength of 0.03 or less, e.g. 0.02 or less, when used at a rate corresponding to 0.4-0.8 g/l surfactant.
- f) An enzymatic liquid detergent composition formulated to
15 give a wash liquor ionic strength of 0.01 or more, e.g. 0.02 or more, when used at a rate corresponding to 0.4-0.8 g/l surfactant.

It was found that the subtilase variants of the present
20 invention can also be usefully incorporated in detergent composition in the form of bars, tablets, sticks and the like for direct application to fabrics, hard surfaces or any other surface. In particular, they can be incorporated into soap or soap/synthetic compositions in bar form, wherein they exhibit a
25 remarkable enzyme stability. Detergent composition in the form of bars, tablets, sticks and the like for direct application, are for example described in South African Patent 93/7274, incorporated herein by reference.

30 Accordingly, the preferred bars in accordance with this invention comprise, in addition to the subtilase variant:

- i) 25 to 80%, most preferably 25 to 70%, by weight of detergent active which is soap or a mixture of soap and synthetic detergent active, reckoned as anhydrous;

SUBSTITUTE SHEET

- ii) 0 to 50 % and, most preferably, 10 to 30% by weight of water;
- iii) 0 to 35% and, most preferably, 0.1 to 30% by weight filler.

5

In general, the amount of subtilase variant to be included in such compositions of the invention is such that it corresponds with a proteolytic activity of 0.1 to 100 GU/mg based on the composition, preferably 0.5 to 20GU/mg, most preferably 1.0 to 10 10 GU/mg, where GU/mg is glycine unit per milligram.

METHOD FOR PRODUCING MUTATIONS IN SUBTILASE GENES

Many methods for introducing mutations into genes are well known in the art. After a brief discussion of cloning subtilase genes, methods for generating mutations in both random sites, and specific sites, within the subtilase gene will be discussed.

20 CLONING A SUBTILASE GENE

The gene encoding a subtilase may be cloned from any of the organisms indicated in Table I, especially gram-positive bacteria or fungus, by various methods, well known in the art. First a genomic, and/or cDNA library of DNA must be constructed using chromosomal DNA or messenger RNA from the organism that produces the subtilase to be studied. Then, if the amino-acid sequence of the subtilase is known, homologous, labelled oligonucleotide probes may be synthesized and used to identify subtilisin-encoding clones from a genomic library of bacterial DNA, or from a cDNA library. Alternatively, a labelled oligonucleotide probe containing sequences homologous to subtilase from another strain of bacteria or organism could be used as a probe to identify subtilase-encoding clones, using hybridization and washing conditions of lower stringency.

35

SUBSTITUTE SHEET

Yet another method for identifying subtilase-producing clones would involve inserting fragments of genomic DNA into an expression vector, such as a plasmid, transforming protease-negative bacteria with the resulting genomic DNA library, and then plating the transformed bacteria onto agar containing a substrate for subtilase, such as skim milk. Those bacteria containing subtilase-bearing plasmid will produce colonies surrounded by a halo of clear agar, due to digestion of the skim milk by excreted subtilase.

10

GENERATION OF RANDOM MUTATIONS IN THE SUBTILASE GENE

Once the subtilase gene has been cloned into a suitable vector, such as a plasmid, several methods can be used to introduce random mutations into the gene.

15

One method would be to incorporate the cloned subtilase gene, as part of a retrievable vector, into a mutator strain of *Eschericia coli*.

- 20 Another method would involve generating a single stranded form of the subtilase gene, and then annealing the fragment of DNA containing the subtilase gene with another DNA fragment such that a portion of the subtilase gene remained single stranded. This discrete, single stranded region could then be exposed to
- 25 any of a number of mutagenizing agents, including, but not limited to, sodium bisulfite, hydroxylamine, nitrous acid, formic acid, or hydralazine. A specific example of this method for generating random mutations is described by Shortle and Nathans (1978, *Proc. Natl. Acad. Sci. U.S.A.*, 75 2170-2174).
- 30 According to the shortle and Nathans method, the plasmid bearing the subtilase gene would be nicked by a restriction enzyme that cleaves within the gene. This nick would be widened into a gap using the exonuclease action of DNA polymerase I. The resulting single-stranded gap could then be mutagenized
- 35 using any one of the above mentioned mutagenizing agents.

SUBSTITUTE SHEET

Alternatively, the subtilisin gene from a *Bacillus* species including the natural promoter and other control sequences could be cloned into a plasmid vector containing replicons for both *E. coli* and *B. subtilis*, a selectable phenotypic marker and the M13 origin of replication for production of single-stranded plasmid DNA upon superinfection with helper phage IR1. Single-stranded plasmid DNA containing the cloned subtilisin gene is isolated and annealed with a DNA fragment containing vector sequences but not the coding region of subtilisin, resulting in a gapped duplex molecule. Mutations are introduced into the subtilisin gene either with sodium bisulfite, nitrous acid or formic acid or by replication in a mutator strain of *E. coli* as described above. Since sodium bisulfite reacts exclusively with cytosine in a single-stranded DNA, the mutations created with this mutagen are restricted only to the coding regions. Reaction time and bisulfite concentration are varied in different experiments such that from one to five mutations are created per subtilisin gene on average. Incubation of 10 mg of gapped duplex DNA in 4 M Na-bisulfite, pH. 6.0, for 9 minutes at 37°C in a reaction volume of 400 ml, deaminates about 1% of cytosines in the single-stranded region. The coding region of mature subtilisin contains about 200 cytosines, depending on the DNA strand. Advantageously, the reaction time is varied from about 4 minutes (to produce a mutation frequency of about one in 200) to about 20 minutes (about 5 in 200).

After mutagenesis the gapped molecules are treated *in vitro* with DNA polymerase I (Klenow fragment) to make fully double-stranded molecules and fix the mutations. Competent *E. coli* are then transformed with the mutagenized DNA to produce an amplified library of mutant subtilisins. Amplified mutant libraries can also be made by growing the plasmid DNA in a Mut

SUBSTITUTE SHEET

D strain of *E. coli* which increases the range of mutations due to its error prone DNA polymerase.

The mutagens nitrous acid and formic acid may also be used to produce mutant libraries. Because these chemicals are not as specific for single-stranded DNA as sodium bisulfite, the mutagenesis reactions are performed according to the following procedure. The coding portion of the subtilisin gene is cloned in M13 phage by standard methods and single stranded phage DNA prepared. The single-stranded DNA is then reacted with 1 M nitrous acid pH. 4.3 for 15-60 minutes at 23°C or 2.4 M formic acid for 1-5 minutes at 23°C. These ranges of reaction times produce a mutation frequency of from 1 in 1000 to 5 in 1000. After mutagenesis, a universal primer is annealed to the M13 DNA and duplex DNA is synthesized using the mutagenized single-stranded DNA as a template so that the coding portion of the subtilisin gene becomes fully double-stranded. At this point the coding region can be cut out of the M13 vector with restriction enzymes and ligated into an un-mutagenized expression vector so that mutations occur only in the restriction fragment. (Myers et al., *Science* 229 242-257 (1985)).

GENERATION OF SITE DIRECTED MUTATIONS IN THE SUBTILASE GENE

Once the subtilase gene has been cloned, and desirable sites for mutation identified and the residue to substitute for the original ones have been decided, these mutations can be introduced using synthetic oligonucleotides. These oligonucleotides contain nucleotide sequences flanking the desired mutation sites; mutant nucleotides are inserted during oligonucleotide synthesis. In a preferred method, a single stranded gap of DNA, bridging the subtilase gene, is created in a vector bearing the subtilase gene. Then the synthetic nucleotide, bearing the desired mutation, is annealed to a homologous portion of the single-stranded DNA. The remaining gap is then

SUBSTITUTE SHEET

filled in by DNA polymerase I (Klenow fragment) and the construct is ligated using T4 ligase. A specific example of this method is described in Morinaga et al., (1984, *Biotechnology* 2 646-639). According to Morinaga et al., a fragment within the gene is removed using restriction endonuclease. The vector/gene, now containing a gap, is then denatured and hybridized to a vector/gene which, instead of containing a gap, has been cleaved with another restriction endonuclease at a site outside the area involved in the gap. A single-stranded region of the gene is then available for hybridization with mutated oligonucleotides, the remaining gap is filled in by the Klenow fragment of DNA polymerase I, the insertions are ligated with T4 DNA ligase, and, after one cycle of replication, a double-stranded plasmid bearing the desired mutation is produced. The Morinaga method obviates the additional manipulation of constructing new restriction sites, and therefore facilitates the generation of mutations at multiple sites. U.S. Reissue Patent number 34,606 by Estell et al., issued May 10, 1994, is able to introduce oligonucleotides bearing multiple mutations by performing minor alterations of the cassette, however, an even greater variety of mutations can be introduced at any one time by the Morinaga method, because a multitude of oligonucleotides, of various lengths, can be introduced.

25

EXPRESSION OF SUBTILASE MUTANTS

According to the invention, a mutated subtilase gene produced by methods described above, or any alternative methods known in the art, can be expressed, in enzyme form, using an expression vector. An expression vector generally falls under the definition of a cloning vector, since an expression vector usually includes the components of a typical cloning vector, namely, an element that permits autonomous replication of the vector in a microorganism independent of the genome of the microorganism, and one or more phenotypic markers for selection

SUBSTITUTE SHEET

purposes. An expression vector includes control sequences encoding a promoter, operator, ribosome binding site, translation initiation signal, and, optionally, a repressor gene or various activator genes. To permit the secretion of the expressed protein, nucleotides encoding a "signal sequence" may be inserted prior to the coding sequence of the gene. For expression under the direction of control sequences, a target gene to be treated according to the invention is operably linked to the control sequences in the proper reading frame.

10 Promoter sequences that can be incorporated into plasmid vectors, and which can support the transcription of the mutant subtilase gene, include but are not limited to the prokaryotic β -lactamase promoter (Villa-Komaroff, et al. (1978) *Proc. Natl. Acad. Sci. U.S.A.* 75 3727-3731) and the tac promoter (DeBoer, 15 et al. (1983) *Proc. Natl. Acad. Sci. U.S.A.* 80 21-25). Further references can also be found in "Useful proteins from recombinant bacteria" in *Scientific American* (1980) 242 74-94.

According to one embodiment *B. subtilis* is transformed by an expression vector carrying the mutated DNA. If expression is to take place in a secreting microorganism such as *B. subtilis* a signal sequence may follow the translation initiation signal and precede the DNA sequence of interest. The signal sequence acts to transport the expression product to the cell wall where 25 it is cleaved from the product upon secretion. The term "control sequences" as defined above is intended to include a signal sequence, when it is present.

Other host systems known to the skilled person are also contemplated for the expression and production of the protease 30 variants of the invention. Such host systems comprise fungi, including filamentous fungi, plant, avian and mammalian cells, as well as others.

SUBSTITUTE SHEET

MATERIALS AND METHODS

Strains:

- 5 *B. subtilis* 309 and 147 are variants of *Bacillus lentus*, deposited with the NCIB and accorded the accession numbers NCIB 10309 and 10147, and described in US Patent No. 3,723,250 incorporated by reference herein.
- 10 *E. coli* MC 1000 (M.J. Casadaban and S.N. Cohen (1980); *J. Mol. Biol.* 138 179-207), was made r⁻,m⁺ by conventional methods and is also described in US Patent Application Serial No. 039,298.

Proteolytic Activity

- 15 In the context of this invention proteolytic activity is expressed in Kilo NOVO Protease Units (KNPU). The activity is determined relatively to an enzyme standard (SAVINASE[®]), and the determination is based on the digestion of a dimethyl casein (DMC) solution by the proteolytic enzyme at standard
- 20 conditions, i.e. 50°C, pH 8.3, 9 min. reaction time, 3 min. measuring time. A folder AF 220/1 is available upon request to Novo Nordisk A/S, Denmark, which folder is hereby included by reference.
- 25 A GU is a Glycine Unit, defined as the proteolytic enzyme activity which, under standard conditions, during a 15-minutes' incubation at 40 deg C, with N-acetyl casein as substrate, produces an amount of NH₂-group equivalent to 1 mmole of glycine.
- 30 Enzyme activity can also be measured using the PNA assay, according to reaction with the soluble substrate succinyl-alanine-alanine-proline-phenyl-alanine-para-nitrophenol, which is described in the Journal of American Oil Chemists Society,

SUBSTITUTE SHEET

Rothgeb, T.M., Goodlander, B.D., Garrison, P.H., and Smith, L.A., (1988).

5

EXAMPLES

For the generation of enzyme variants according to the invention the same materials and methods as described in i.a. WO 89/06279 (Novo Nordisk A/S), EP 130,756 (Genentech), EP 10 479,870 (Novo Nordisk A/S), EP 214,435 (Henkel), WO 87/04461 (Amgen), WO 87/05050 (Genex), EP application no. 87303761 (Genentech), EP 260,105 (Genencor), WO 88/06624 (Gist-Brocades NV), WO 88/07578 (Genentech), WO 88/08028 (Genex), WO 88/08033 (Amgen), WO 88/08164 (Genex), Thomas et al. (1985) *Nature*, 318 15 375-376; Thomas et al. (1987) *J. Mol. Biol.*, 193, 803-813; Russel and Fersht (1987) *Nature* 328 496-500. Other methods well established in the art may also be used.

20

EXAMPLE 1

Construction and Expression of Enzyme Variants:

A vector suited to a synthetic gene coding for subtilase 309 and its mutants was constructed. It is essentially a pUC19 25 plasmid [Yanish-Perron and Messing (1985) *Gene*; 33 103-119], in which the multiple cloning site has been replaced by a linker containing the restriction sites used to separate five sub--fragments constituting the gene. The new linker was inserted into EcoRI - HindIII cut pUC19 thereby destroying these sites. 30 The details of this construction are described in WO 92/19729 on pages 25-26 and in figure 1 (sheets 1/7-7/7) thereof, the content of which is hereby included by reference.

Each subfragment was made from 6 to 12 oligonucleotides. The 35 oligonucleotides were synthesised on an automatic DNA

SUBSTITUTE SHEET

synthesizer using phosphoramidite chemistry on a controlled glass support [Beaucage and Carruthers (1981); *Tetrahedron Letters* 22 1859-1869].

5 The five subfragments were isolated on a 2% agarose gel and inserted into pSX191. The sequence was verified by dideoxynucleotide sequencing. Fragments A-E were isolated and ligated together with KpnI-BamHI cut pSX191. The ligation mixtures were used to transform competent *E. coli* MC1000 r^{-}, m^{-} selecting for
10 ampicillin resistance. The 850 bp KpnI-BamHI fragment that constitutes the part of the subtilisin 309 gene coding for the mature part of the enzyme was then used to replace the wild type gene on pSX212 giving rise to pSX222, which was then transformed into a competent *B. subtilis* strain. After
15 fermentation of the transformed strain and purification of the enzyme it was shown that the product was indistinguishable from the wild type product.

Protease variants derived from the synthetic gene are made by
20 using oligonucleotides with altered sequence at the place(s) where mutation is wanted (e.g. with sequences as given below) and mixing them with the rest of the oligonucleotides appropriate to the synthetic gene. Assembly of the variant gene is carried out with the variant materials in a manner otherwise
25 analogous to that described above. Further information on synthetic genes generally is available in Agarwal et al (1970); *Nature*; 227 27-34.

A KpnI site was introduced into the beginning of the subtilase
30 309 synthetic gene encoding the mature part of the enzyme. The method used is called oligonucleotide directed double-strand break repair mutagenesis and is described by Mandecki (1986) *Proc. Nat. Acad. Sci. USA* 83 7177-7181. pSX172 is opened with NcoI at the beginning of the mature part of the subtilase 309
35 gene and is mixed with the oligonucleotide NOR 789 (see WO

SUBSTITUTE SHEET

92/19729), heated to 100°C, cooled to 0°C, and transformed into *E. coli*. After retransformation, the recombinants can be screened by colony hybridisation using 32-P-labelled NOR 789. The recombinants that turned out to be positive during the 5 screening had the KpnI site introduced right in front of NcoI by changing two bases without changing the amino acid sequence. pSX172 is described in EP Patent Publication No. 405 901. The KpnI site so created is inserted into pSX120 on a 400-bp PvuI-NheI fragment, giving rise to pSX212. pSX120 is also 10 described in EP Patent Publication No. 405 901.

The synthetic gene is inserted between KpnI and BamHI on pSX212, giving rise to pSX222.

15 Examples of mutations and corresponding sequences of oligonucleotides are as follows:

R170L (fragment D1)

20

```
5'- AATTCAGGTGCAGGCTCAATCAGCTATCCGGCGCTCTAT - 3'
      ||||||||||||||||||||||||||||*|||
5'-  GTCCACGTCCGAGTTAGTCGATAGGCCGCGAGATACGCTTG - 3'
```

25 R170I (fragment D1)

```
5'- AATTCAGGTGCAGGCTCAATCAGCTATCCGGCGATCTAT - 3'
      ||||||||||||||||||||||||||||**|||
5'-  GTCCACGTCCGAGTTAGTCGATAGGCCGCTAGATACGCTTG - 3'
```

30

S57P (fragment B1)

```
5' - AGCTTTGTACCAGGGGAACCGCCGACTCAAGATGGG - 3'
      ||||||||||||||||||*|||||||
35 3' - AACATGGTCCCCTTGGCGGCTGAGTTCTACCCCTTACCC - 5'
```

These oligonucleotides were combined with the rest of the oligonucleotides from the synthetic gene that was not changed.

SUBSTITUTE SHEET

EXAMPLE 2

Purification of Enzyme Variants:

5 This procedure relates to purification of a 10 litre scale fermentation of the Subtilisin 147 enzyme, the Subtilisin 309 enzyme or mutants thereof.

Approximately 8 litres of fermentation broth were centrifuged
10 at 5000 rpm for 35 minutes in 1 litre beakers. The supernatants were adjusted to pH 6.5 using 10% acetic acid and filtered on Seitz Supra S100 filter plates.

The filtrates were concentrated to approximately 400 ml using
15 an Amicon CH2A UF unit equipped with an Amicon S1Y10 UF cartridge. The UF concentrate was centrifuged and filtered prior to absorption at room temperature on a Bacitracin affinity column at pH 7. The protease was eluted from the Bacitracin column at room temperature using 25% 2-propanol and
20 1 M sodium chloride in a buffer solution with 0.01 dimethylglutaric acid, 0.1 M boric acid and 0.002 M calcium chloride adjusted to pH 7.

The fractions with protease activity from the Bacitracin
25 purification step were combined and applied to a 750 ml Sephadex G25 column (5 cm dia.) equilibrated with a buffer containing 0.01 dimethylglutaric acid, 0.2 M boric acid and 0.002 m calcium chloride adjusted to pH 6.5.

30 Fractions with proteolytic activity from the Sephadex G25 column were combined and applied to a 150 ml CM Sepharose CL 6B cation exchange column (5 cm dia.) equilibrated with a buffer containing 0.01 M dimethylglutaric acid, 0.2 M boric acid, and 0.002 M calcium chloride adjusted to pH 6.5.

35

SUBSTITUTE SHEET

The protease was eluted using a linear gradient of 0-0.1 M sodium chloride in 2 litres of the same buffer (0-0.2 M sodium chloride in case of Subtilisin 147).

5 In a final purification step protease containing fractions from the CM Sepharose column were combined and concentrated in an Amicon ultrafiltration cell equipped with a GR81PP membrane (from the Danish Sugar Factories Inc.).

10 By using the techniques of Example 1 for the construction and the above isolation procedure the following subtilisin 309 variants were produced and isolated:

A:	G159I
15 B:	S164I
C:	Y167I
D:	R170I
E:	R170L
F:	R170M
20 G:	R170F
H:	G195F
I:	S57P+R170L
J:	R170L+N218S
K:	S57P+R170L+N218S
25 L:	R170L+N218S+M222A
M:	S57P+R170L+S188P+A194P
N:	Y167I+R170L
O:	S57P+R170L+Q206E
P:	R170L+Q206E
30 Q:	Y167I+R170L+Q206E
R:	Y167I+R170L+A194P
S:	Y167I+R170L+N218S
T:	Y167I+R170L+A194P+N218S
U:	Y167I+Y171I
35 V:	R170G

SUBSTITUTE SHEET

W: R170C
X: Y171I
Y: Y167I+R170L+N218S

5

EXAMPLE 3Stability in Detergent Compositions Comprising Enzyme Variants**10 Example D1:**

An (isotropic) aqueous detergent liquid according to an embodiment of the invention is formulated to contain:

SUBSTITUTE SHEET

<u>Ingredient</u>	<u>%</u>
NaLAS	8.0
Neodol 25-9	8.0
AES 25-3S	14.0
NaCitrate.2H ₂ O	5.0
Propylene Glycol	5.0
Sorbitol	4.5
F-dye Tinopal UNPA-GX	0.15
Lytron 614 Opacifier	0.03
Kathon Preservative	0.0003
Acid Blue 80	0.00117
Acid Violet 48	0.0033
SAVINASE® 16L	0.25
LIPOLASE® 100L	0.70
Fragrance	0.15
Water	ad 100.0

The pH is adjusted to 7.1.

5 Table III.

Residual enzyme activity (in percentage of original activity) after storage at 37°C for Example D1 comprising the BLS309 variant S57P+R170L+N218S.

SUBSTITUTE SHEET

Storage time (days)	Wild-type	S57P+R170L+N218S
0	100	100
3	44	74
7	11	50
5 10	5	36
14	7	27

From Table III it is evident that the variant S57P+R170L+N218S exhibits a remarkably improved stability in this type of detergent. Moreover, the variant S57P+R170L+N218S possesses excellent compatibility towards lipase.

Table IV.

Residual lipase activity (in percentage of original activity) after storage at 37°C for Example D1 comprising the BLS309 variant S57P+R170L+N218S and LIPOLASE®.

Storage time (days)

20 LIPOLASE® plus:	Wild-type	S57P+R170L+N218S
0	100	100
3	38	67
7	24	44
10	22	33
25 14	21	27

From the above Table IV it is apparent that, in addition to the stability of the protease, the compatibility of the protease is also improved.

30

SUBSTITUTE SHEET

Example D2:

A non-aqueous detergent liquid according to an embodiment of the invention is formulated using 38.5% C13-C15 linear primary
 5 alcohol alkoxylated with 4.9 mol/mol ethylene oxide and 2.7 mol/mol propylene oxide, 5% triacetin, 30% sodium triphosphate, 4% soda ash, 15.5% sodium perborate monohydrate containing a minor proportion of oxoborate, 4% TAED, 0.25% EDTA of which 0.1% as phosphonic acid, Aerosil 0.6%, SCMC 1%, and 0.6%
 10 protease. The pH is adjusted to a value between 9 and 10, e.g. about 9.8.

Example D3:

Structured liquid detergents can for example contain, in
 15 addition to a protease as described herein, 2-15% nonionic surfactant, 5-40% total surfactant, comprising nonionic and optionally anionic surfactant, 5-35% phosphate-containing or non-phosphate containing builder, 0.2-0.8% polymeric thickener, e.g. cross-linked acrylic polymer with m.w. over 10^6 , at least
 20 10% sodium silicate, e.g. as neutral waterglass, alkali (e.g. potassium-containing alkali) to adjust to desired pH, preferably in the range 9-10 or upwards, e.g. above pH 11, with a ratio sodium cation: silicate anion (as free silica) (by weight) less than 0.7:1, and viscosity of 0.3-30 Pas (at 20°C
 25 and 20^{s-1}).

Suitable examples contain about 5% nonionic surfactant C13-15 alcohol alkoxylated with about 5 EO groups per mole and with about 2.7 PO groups per mole, 15-23% neutral waterglass with
 30 3.5 weight ratio between silica and sodium oxide, 13-19% KOH, 8-23% STPP, 0-11% sodium carbonate, 0.5% Carbopol 941 (TM).

Protease may be incorporated at for example 0.5%.

Example D4:

(Decoupling polymer liquid)

Priolene 6907	4.5
KOH	10
5 Ethoxylated Alcohol.7EO (Synperonic A7)	4.5
Ethoxylated Alcohol.3EO (Synperonic A3)	4.5
Zeolite 4A	15
Fluorescer Tinopal CBS-X	0.08
Narlex DC1	1
10 Citric acid	8.23
Antifoam silicone DB100	0.3
LAS acid	16.5
Perfume	0.5
Water to	100

15

Table V.

Residual enzyme activity (in percentage of original activity) after storage at 37°C for Example D4 comprising the R170L variant of BLS309.

20

Storage time (days)	R170L	Wild-type
0	100	100
2	98	73
4	96	66
25 10	94	46
33	87	8
81	78	2.1
101	71	0

30 From Table V it is evident that the R170L variant exhibits a remarkably improved stability in this type of detergent.

SUBSTITUTE SHEET

Table VI

Storage (days)	Enzy- me	WT	R170M	S57P+R170L +Q206E	Y167I+R170L+ N218S
0		100	100	100	100
0.1		90.2	78	97	94
1		58	53	95	68
2		40	34	87	55
5		16	27	75	29
6		12	22	73	24
8		8	19	77	17
14		2	11	52	4

From Table VI it can be seen that the variants tested exhibit improved stability in comparison to the wild type enzyme in this type of detergent

5

SUBSTITUTE SHEET

Example D5:

(Decoupling polymer liquid)

Priolene 6907	4.5
KOH	10
5 Ethoxylated Alcohol.7EO (Synperonic A7)	4.5
Ethoxylated Alcohol.3EO (Synperonic A3)	4.5
Zeolite 4A	15
Fluorescer Tinopal CBS-X	0.08
Narlex DC1	1
10 Citric acid	8.23
Antifoam silicone DB100	0.3
LAS acid	16.5
Lipolase® 100L	0.6
Perfume	0.5
15 Water to	100

Table VII.

Residual enzyme activity (in percentage of original activity)
after storage at 37°C for Example D5 comprising the BLS309
20 variant S57P+R170L+N218S.

Storage time (days)	Residual protease activity		Residual lipase activity	
	S57P+R170L+N218S	Wild-type	R170L	S57P+R170L+N218S
25 0	100	100	100	100
2	-	27	41	94
5	97	9	15	76
8	87	4	7	71
12	91	2.4	12	78
30 28	100	2.4	12	70

From Table VII it is evident that the variant S57P+R170L+N218S
exhibits a remarkably improved stability in this type of
detergent. Moreover the variant S57P+R170L+N218S possesses
35 excellent compatibility towards lipase.

SUBSTITUTE SHEET

Example D6:

Soap bars were produced containing 49.7 wt. 80/20 tallow 5 /coconut soap, 49.0% water, 20% sodium citrate, 1.0% citric acid and 0.031% protease. After preparation of the soap bars they were stored at ambient temperature and after specific time intervals samples were taken and measured for protease activity. The stability data are given below:

10

Table VIII

Storage (days)	Enzy-me	WT	R170L	R170L+N218S+S57P	R170L+Y167I
0		100	100	100	100
1		50	100	97	94
2		25	91	100	83
3		-	100	94	80
6		-	98	89	90
10		0	100	94	71
17		-	93	80	73
27		-	95	86	70

From the above Table it is evident that the subtilase variants R170L, R170L+N218S+S57P and R170L+Y167I exhibit a remarkably 15 improved stability in this type of detergent.

Example D7:

Soap bars were produced containing 63.88% 80/20 tallow/coconut soap, 1% coconut fatty acid, 25.1% water, 10% sodium citrate 20 and 0.021% protease. The laundry soap bars were stored at 37°C

SUBSTITUTE SHEET

and after specific time intervals samples were taken and measured for protease activity.

Table IX, Stability data:

Storage (days)	Enzyme	WT	R170L+N218S+S57P
0		100	100
10		10	90.1
14		-	81.5
20		0	91.4
31		-	72.8
35		-	79
45		-	78

5

From the above Table it is evident that the subtilase variant R170L+N218S+S57P exhibits a remarkably improved stability in this type of detergent.

10

EXAMPLE 4

Wash Performance of Detergent Compositions Comprising Enzyme Variants

15

The following examples provide results from a number of washing tests that were conducted under the conditions indicated

SUBSTITUTE SHEET

Experimental conditions

Table X: Experimental conditions for evaluation of Subtilisin 309 variants.

5

Detergent	Protease model detergent '95
Detergent dose	3 g/l
pH	9.5
Wash time	15 min.
Temperature	15°C
Water hardness	9°dH ~ 1.61 mM Ca ²⁺ /Mg ²⁺
Enzymes	Subtilisin 309 variants as listed below
Enzyme conc.	0; 0.1, 0.2, 0.3, 0.4, 0.5, 1.0, 2.0, 3.0 mg/l
Test system	150 ml beakers with a stirring rod.
Cloth/volume	5 cloths (Ø 2.5 cm) / 50 ml Detergent solution.
Cloth	Cotton soiled with grass juice

Subsequent to washing the cloths were flushed in tap water and air-dried.

- 10 The above model detergent is a simple detergent formulation. The most characteristic features are that STP is used as builder and the content of anionic tenside (LAS) is quite high. Further the pH is adjusted to 9.5, which is low for a powder detergent.

SUBSTITUTE SHEET

Table XI

The composition of the model detergent is as follows:

25 %	STP ($\text{Na}_5\text{P}_3\text{O}_{10}$)
25 %	Na_2SO_4
5 10 %	Na_2CO_3
20 %	LAS (Nansa 80S)
5 %	NI (Dobanol 25-7)
5 %	$\text{Na}_2\text{Si}_2\text{O}_5$
0.5 %	Carboxymethylcellulose (CMC)
10 9.5 %	water

dose: 3 g/l

pH is adjusted to 9.5

15 Measurement of remission (R) on the test material has been done at 460 nm using an Elrepho 2000 photometer (without UV). The measured values have been fitted to the expression:

$$\Delta R = \frac{a \cdot \Delta R_{\max} \cdot c}{\Delta R_{\max} + a \cdot c}$$

20 The improvement factor is calculated by use of the initial

slope of the curve: $IF = \frac{a}{a_{ref}}$

DR is the wash effect of the enzyme in remission units.

a is the initial slope of the fitted curve ($c \rightarrow 0$).

a_{ref} is the initial slope for the reference enzyme.

25 c is the enzyme concentration in mg/l

DR_{\max} is the theoretical maximum wash effect of the enzyme in remission units ($c \rightarrow \infty$).

SUBSTITUTE SHEET

Table XII: Variants and improvement factors for Subtilisin 309.

Designation	Variant	IF
S003*	R170Y	2.8
S004*	R170Y + G195E	2.6
S012*	R170Y + G195E + K251E	1.6
G	R170F	3.3
E	R170L	3.8
F	R170M	2.4
D	R170I	4.1
I	S57P + R170L	3.9
J	R170L + N218S	1.6
K	S57P + R170L + N218S	2.3
N	Y167I + R170L	6.2
P	R170L + Q206E	2.6
V	R170G	2.0
W	R170C	3.4
O	S57P + R170L + Q206E	2.9
Q	Y167I + R170L + Q206E	2.4
R	Y167I + R170L + A194P	5.1
X	Y171I	1.2
Y	Y167I + R170L + N218S	4.0
T	Y167I + R170I + A194P + N218S	3.6

* Described in WO 91/00345

5 As it can be seen from Table XII all the Subtilisin 309 variants of the invention exhibits an improvement in wash performance.

SUBSTITUTE SHEET

Table XIII: Variants and improvement factors for Subtilisin 309 in a detergent as described in Example D4

Designation	Variant	IF
S003*	R170Y	1.5
F	R170M	1.2
O	S57P + R170L + Q206E	5.0
X	Y171I	4.2
R	Y167I + R170L + A194P	1.2
T	Y167I + R170L + A194P + N218S	2.0
Y	Y167I + R170L + N218S	2.3

5 *Described in WO 91/00345

As it can be seen from Table XIII all the Subtilisin 309 variants of the invention exhibits an improvement in wash performance.

10

SUBSTITUTE SHEET

PATENT CLAIMS

1. A subtilase variant having improved storage stability and/or improved performance in detergents, wherein one or more
5 amino acid residues situated in, or in the vicinity of a hydrophobic domain of the parent subtilase have been substituted for an amino acid residue more hydrophobic than the original residue, said hydrophobic domain comprising the residues corresponding to residues P129, P131, I165, Y167, Y171
10 of BLS309 (in BASBPN numbering), and said residues in the vicinity thereof comprises residues corresponding to the residues E136, G159, S164, R170, A194, and G195 of BLS309 (in BASBPN numbering), with the exception of the R170M, R170I and R170V variants of BABP92 and the R170Y variant of BLS309.

15

2. The variant of claim 1, wherein said variant exhibits improved stability in liquid detergents.

3. The variant of claim 1, wherein said variant exhibits
20 improved stability in detergents in shaped solid form.

4. The variant of claim 1, wherein said variant exhibits improved wash performance.

25 5. The variant of any of the claims 1 to 4, wherein the original amino acid residue has been substituted for any other residue which increase the hydrophobicity of the original residue, where the substituted residue preferably is a residue selected from the group comprising Val (V), Ile (I), Leu (L),
30 Met (M), Phe (F), Pro (P), and Trp(W), especially Val, Ile or Leu.

6. The variant of any of the claims 1 to 5, wherein the parent subtilase is chosen from the sub-group I-S1.

35

SUBSTITUTE SHEET

7. The variant of claim 6, wherein the parent subtilase is chosen from the group comprising ABSS168, BASBPN, BSSDY, and BLSCAR.
- 5 8. The variant of any of the claims 1 to 5, wherein the parent subtilase is chosen from the sub-group I-S2.
9. The variant of claim 8, wherein the parent subtilase is chosen from the group comprising BLS147, BLS309, BAPB92 AND
10 BYSYAB.
10. The variant of claim 8, wherein the parent subtilase is TVTHER.
- 15 11. The variant of any of claims 1 to 10, wherein said substitution(s) is/are combined with substitutions, insertions or deletions in any other position.
12. The variant of claim 11, wherein said
20 substitution(s) is/are combined with substitutions, insertions or deletions in any of the positions 36, 222, 218, 76.
13. The variant of any of claims 1 to 12, being any of the following:
- 25
- T129V, T129I, T129L, T129M, T129F
A129V, A129I, A129L, A129M, A129F
G131V, G131I, G131L, G131M, G131F
K136V, K136I, K136L, K136M, K136F,
30 S159V, S159I, S159L, S159M, S159F,
T164V, T164I, T164L, T164M, T164F,
K170V, K170I, K170L, K170M, K170F,
Q136V, Q136I, Q136L, Q136M, Q136F,
T159V, T159I, T159L, T159M, T159F,
35 A164V, A164I, A164L, A164M, A164F,

SUBSTITUTE SHEET

Y170V, Y170I, Y170L, Y170M, Y170F,
 Y171A, Y171H, Y171N, Y171P, Y171C, Y171W,
 Y171Q, Y171S, Y171T, Y171G
 Y171V, Y171I, Y171L, Y171M, Y171F
 5 S194V, S194I, S194L, S194M, S194F,
 E136V, E136I, E136L, E136M, E136F,
 G159V, G159I, G159L, G159M, G159F,
 G164V, G164I, G164L, G164M, G164F,
 S164V, S164I, S164L, S164M, S164F,
 10 Y167A, Y167H, Y167N, Y167P, Y167C, Y167W,
 Y167Q, Y167S, Y167T, Y167G
 Y167V, Y167I, Y167L, Y167M, Y167F
 R170A, R170H, R170N, R170P, R170C, R170W
 R170Q, R170S, R170T, R170Y, R170G
 15 R170V, R170I, R170L, R170M, R170F,
 A194V, A194I, A194L, A194M, A194F,
 P194V, P194I, P194L, P194M, P194F,
 E195V, E195I, E195L, E195M, E195F,
 G195V, G195I, G195L, G195M, G195F,

20

14. The variant of any of claims 1 to 13, wherein said
 variant is combined with further substitutions, deletions
 and/or insertions in any one or more of the positions:

27, 36, 57, 76, 97, 101, 104, 120, 123, 206, 218, 222, 224, 235
 25 and 274.

15. The variant of claim 14, wherein said subtilase
 belongs to the I-S2 sub-group and said further change is chosen
 from the group comprising K27R, *36D, S57P, N76D, G97N, S101G,
 30 V104A, V104N, V104Y, H120D, N123S, A194P, Q206E, N218S, M222S,
 M222A, T224S, K235L, and T274A.

16. The variant of claim 15 comprising any one or two of
 the substitutions X167V, X167M, X167F, X167L, X167I, X170V,
 35 X170M, X170F, X170L, and/or X170I in combination with any one

SUBSTITUTE SHEET

or more of the other substitutions, deletions and/or insertions mentioned in any of claims 1 to 15.

17. The variant of claim 15 comprising any of the 5 variants V104N+S101G, K27R+V104Y+N123S+T274A, or N76D+V104A, or other combinations of these mutations (V104N, S101G, K27R, V104Y, N123S, T274A, N76D, V104A), in combination with any one or more of the substitutions, deletions and/or insertions mentioned in any of claims 1 to 15.

10

18. The variant of claim 15 or 16, chosen from the group of variants comprising:

- a) S57P+R170L
- 15 a') S57P+R170I
- b) R170L+N218S
- b') R170I+N218S
- c) S57P+R170L+N218S
- c') S57P+R170I+N218S
- 20 c'') S57P+V104Y+R170L+N218S
- c''') S57P+V104Y+R170I+N218S
- d) R170L+N218S+M222A
- d') R170I+N218S+M222S
- d'') R170L+N218S+M222A
- 25 d''') R170I+N218S+M222S
- e) S57P+R170L+S188P+A194P
- e') S57P+R170I+S188P+A194P
- f) Y167L+R170L
- f') Y167L+R170I
- 30 g) Y167I+R170L
- g') Y167I+R170I
- h) N76D+R170L+N218S
- h') N76D+R170I+N218S
- i) S57P+N76D+R170L+N218S
- 35 i') S57P+N76D+R170I+N218S

SUBSTITUTE SHEET

- j) N76D+R170L+N218S+M222A
- j') N76D+R170I+N218S+M222S
- j") N76D+R170L+N218S+M222A
- j''') N76D+R170L+N218S+M222S
- 5 k) S57P+R170I+S188P+A194P+N218S
- k') S57P+R170I+S188P+A194P+N218S
- l) *36D+N76D+H120D+R170L+G195E+K235L
- l') *36D+N76D+H120D+R170I+G195E+K235L
- l") *36D+N76D+H120D+Y167I+R170L+G195E+K235L
- 10 l''') *36D+N76D+H120D+Y167I+R170I+G195E+K235L
- m) N76D+H120D+R170L+G195E+K235L
- m') N76D+H120D+R170I+G195E+K235L
- m") N76D+H120D+Y167I+R170L+G195E+K235L
- m''') N76D+H120D+Y167I+R170I+G195E+K235L
- 15 n) *36D+G97N+V104Y+H120D+R170L+A194P+G195E+K235L
- n') *36D+G97N+V104Y+H120D+R170I+A194P+G195E+K235L
- o) S57P+R170L+Q206E
- o') S57P+R170I+Q206E
- p) R170L+Q206E
- 20 p') R170I+Q206E
- q) Y167I+R170L+Q206E
- q') Y167I+R170I+Q206E
- r) Y167F+R170L
- r') Y167F+R170I
- 25 t) Y167I+R170L+A194P
- t') Y167I+R170I+A194P
- t") Y167L+R170L+A194P
- t''') Y167L+R170I+A194P
- u) Y167I+R170L+N218S
- 30 u') Y167I+R170I+N218S
- u") Y167L+R170L+N218S
- u''') Y167L+R170I+N218S
- v) Y167I+R170L+A194P+N218S
- v') Y167I+R170I+A194P+N218S
- 35 v") Y167L+R170L+A194P+N218S

SUBSTITUTE SHEET

v'') Y167L+R170I+A194P+N218S
 x) R170L+P131V
 x') R170I+P131V
 y) *36D+Y167I+R170L
 5 y') *36D+Y167I+R170I
 z) Y167I+Y171I
 aa) Y167V+R170L
 aa') Y167V+R170I
 bb) R170L+Y171I
 10 bb') R170I+Y171L
 bb'') R170L+Y171L
 bb''') R170I+Y171I
 cc) Y167I+Y171L+N218S
 cc') Y167I+Y171I+N218S

15

19. A process which comprises effecting a mutation in DNA encoding a subtilase enzyme or its pre- or preproenzyme at one or more of the positions corresponding amino acid residues in or in the vicinity of a hydrophobic domain of the parent
 20 subtilase have been substituted for an amino acid residue more hydrophobic than the original residue, said hydrophobic domain comprising the residues P129, P131, I165, Y167, Y171 of BLS309, and the residues in the vicinity thereof for BLS309 are E136, G159, S164, R170, A194, and G195, with the exception of the
 25 R170M, R170I and R170V variants of BABP92, and testing for improved storage stability and/or wash performance in detergents.

20. A process which comprises the manufacture of a mutant
 30 subtilase enzyme having a desired improved storage stability or wash performance following its identification according to the procedure of claim 19.

21. A DNA sequence encoding a subtilase variant of any of the
 35 claims 1 to 18.

SUBSTITUTE SHEET

22. A vector comprising a DNA sequence of claim 21.

23. A microbial host transformed with a vector of claim 22.

5

24. The microbial host of claim 23, which is a bacterium, preferably a *Bacillus*.

25. The microbial host of claim 23, which is a fungus or
10 yeast, preferably a filamentous fungus, especially an *Aspergillus*.

26. A method for producing a variant of any of claims 1 to 18,
wherein a host of any of claims 23 to 25 is cultured under
15 conditions conducive to the expression and secretion of said
variant, and the variant is recovered.

SUBSTITUTE SHEET

1/2

	130	140	160	170	190	200
BASBPNGGPGS	---	SAAL KAAVDKAVAS	EGTSGSS	TVGYPGKYP	FSS	VGPE
BSS168 GGPTG	---	STAL KTVDKAVSS	EGSSGTS	TVGYPKYP	FSS	AGSE
BSSDY GGPGS	---	STAL KQAVDKAVAS	SGSSGQN	TIGYPAKYD	FSS	VGAE
BLSCARGGPGS	---	STAH KQAVDNAYAR	SGSSGNTN	TIGYPAKYD	FSS	VGAE
BAPB92 GSPSP	---	SATLEQAVNSATSR	SCAG	SISYPARYA	FSQ	YGAG
BYSYABGSSAG	---	SATHEQAVNQATAS	SCAG	NVGFARYA	FSQ	YGAG
BLS147 GSTSG	---	SSTLELAVNRANNA	TGRQ	GVNYPARYS	FST	YGPE
BSEPR GTTSD	---	SKILHDAVNKAYEQ	DGNGK	PVNYPAAYS	FST	TGDE
BSISPI GGPSD	---	VPELEAVNKAVKN	EGDGDERT	ELSYPAAYN	FSN	ANKE
TVTHERGGTVG	---	NSGLQAVNYAWNK	AGNT	APNYPAYS	FST	YGSW
ONEBPRGGGG	---	CSQNSQRHIDKTTNL	ENQDA	SRTWPSSCN	FSN	YGAR
XCEPRGGGS	---	CSTTHQAINGAVSR	DASNV	SGSLPANCA	YSN	FGTC
BSBPF GGGSG	---	LDEWY ROMVNAWRAA	TDLFIPGGPG	SIANPANYP	FSL	QGPSYDEIKPEISAP
EFCYLA GSYKN	---	MEIDDERFTVEAF	RKVNYARKN	EKHIPGGLE	YSN	YGSN
SEPIPGNYLI	(9)	RDDEKVDYDAL	QKAINYAKK	TSKKVYDSPANLN	FSN	YGN
SPSCPA GNAAL	---	AYANLPDET	KKAFDYAKSK	HPDYGVGTPAAAD	FSS	WGLTADGNIKPDIAAP
LLSK11 GNSG	---	NQTLDEPE	LAAVQMANES	SGTSGSATEGVNKDYGLQDNEMVSGPSTR	FTS	YGPVSNLSFKPDITAP
SMEXSP GIAPD	(43)	QPVPTGGHSA	NSTLLRAARH	YNNYNIPEAQKSL	SST	SCQQTAS
AVPRCA GPPDG	(10)	KQKVPDPST	RLANDYAINKG	GNEVD	YSD	FGTA
HMPPCJ GPND	---	GKTVEGPGRLA	QKAFEGYVKQG	GGRQGDNC	YAE	KCSS
HSIPC2 GPTDN	---	GKTVDGPRDVT	LQAHADGVNKG	GGSY-DDCN	YDE	SCSS
HSFURI GPEDD	---	GKTVDGPARLA	EEAFRCVVSQ	GGREHDSCH	YSE	ACSS
DMFURI GPDDD	---	GKTVDGPGELA	SRAFIEGTTKG	GGREQDNCN	YSE	KCSS
KLKEX1 GPSDD	---	GKTMQAPDTLV	KKAIKGVTEG	GGHFGDSCN	YSE	SCSA
SCKEX2 GPADD	---	GRHLQGPSDLV	KKALVKGVTEG	GGRQDNCN	YSE	GCSA
VAPROA GGGQS	---	VALDSAVQSAVQS	SNA-DACN	YSPARVA	FSN	WGSC
TRT41A GGGAS	---	TALDTAVHNAINA	DNR-DACF	YSPARVT	FSN	YGR
TAAQUA GGGVS	---	TALDNVKNISIAA	DNA-NACN	YSPARVA	FSN	YGSC
TAPROK GGGYS	---	SSVNSAARLQSS	NNA-DARN	YSPASEP	FSN	YGSV
TAPROR GGGYS	---	SSVNSAANLQSS	NNA-DARN	YSPASEE	FSN	YGSV
TAPROT GGPSS	---	SAVNRAAAEITSA	EAT-DASS	YSPASAA	FSN	YGSV
ACALPR GGGYS	---	SAFNNAVNTAYSR	DNQ-NAAN	YSPASAP	FSN	YGSV
AOALPR GGGYS	---	KAFNDAVENAFEQ	ENS-DAGQ	TSPASAD	FSN	YGSV
SCPRB1 GGGKS	---	PALDLAVNAAVEV	ENQ-DACN	TSPASAD	FSN	YGSV
YLXPR2 GGPKS	---	ASQDALWSRATQE	DAV-DACN	DSPGNIGGWSGGQSNYGT	---	YGSV

FIG. 1

SUBSTITUTE SHEET




FIG. 2

SUBSTITUTE SHEET

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C12N 9/54

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI, EPOQUE, US PAT FULL, MEDLINE, BIOSIS, DERWENT BIOTECH ABSTRACTS, EMBL/GENBANK/DBBJ/SWISSPROT (STRAND), REGISTRY FILE, CA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0525610 A2 (SOLVAY ENZYMES GMBH & CO. KG), 3 February 1993 (03.02.93), page 3, line 154, claim 7, figure 1, figure 3 --	1-26
X	EP 0405901 A1 (UNILEVER PLC), 2 January 1991 (02.01.91), claim 1 --	1-26
X	WO 8906279 A1 (NOVO INDUSTRI A/S), 13 July 1989 (13.07.89), claim 1 --	1-26
X	WO 9100345 A1 (NOVO NORDISK A/S), 10 January 1991 (10.01.91), claim 1 -----	1-26

☐ Further documents are listed in the continuation of Box C.
 ☒ See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

5 August 1996

Date of mailing of the international search report

08 -08- 1996

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Patrick Andersson
Telephone No. +46 8 782 25 00

Form PCT/ISA/210 (second sheet) (July 1992)

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
EP-A2-	0525610	03/02/93	DE-A,A-	4224125	28/01/93
			US-A-	5453372	26/09/95
EP-A1-	0405901	02/01/91	CA-A-	2034486	27/12/90
			JP-T-	4500385	23/01/92
			WO-A,A-	9100334	10/01/91
			EP-A,A-	0405902	02/01/91
			JP-T-	4500384	23/01/92
			WO-A,A-	9100335	10/01/91
WO-A1-	8906279	13/07/89	AT-T-	136329	15/04/96
			DE-D-	68926163	00/00/00
			EP-A,A,B	0396608	14/11/90
			EP-A,A-	0675196	04/10/95
			JP-T-	3503477	08/08/91
			JP-B-	6075504	28/09/94
			JP-A-	6292577	21/10/94
WO-A1-	9100345	10/01/91	AU-A-	5956690	17/01/91
			CA-A-	2062732	27/12/90
			EP-A,A-	0479870	15/04/92
			JP-T-	4505856	15/10/92

Form PCT/ISA/210 (patent family annex) (July 1992)